

## Diabetes Reversal by Plant-Based Diet

Biswaroop Roy Chowdhury

Medical Nutritionist, Indo-Vietnam Medical Board, India

### Corresponding author

Dr. Biswaroop Roy Chowdhury, Medical Nutritionist, Indo-Vietnam Medical Board, India. E-mail: biswaroop@biswaroop.com.

Submitted: 17 Aug 2017; Accepted: 24 Aug 2017; Published: 05 Dec 2017

### Abstract

**Background:** Diabetes causes a never-ending medicine and or insulin treatment for the diseased. Also, the patients are bound to follow a particular diet, with eliminating most of the sugary foods; that further deteriorates the quality of life. This gave way to the study, focused on diabetes cure without medicines and on rich fruits and vegetable diet.

The clinical trial on 55 diabetes patients with a team of 6 medical associates was practiced for 3-days at Zorba, The Buddha, 10 - Tropical Drive, Ghitorni, New Delhi from 29th April to 1st May, 2016. The goal was to establish and observe the effects of plant-based diet on the sugar levels of the diabetes patients. These included both insulin-dependent and independent, diabetes type-1 and type-2 patients. The 3-days Residential Treatment Tour involved 55 subjects with different age groups and demographic profiles. The study considered participants from different countries to find out the global impact of the treatment.

### Objective

**Aim:** The burden of the disease diabetes is rising globally. The aim of the research is to find out that on discontinuing the medicines and being on a particular plant-based diet, can high blood glucose levels in diabetes patients be normalized.

### Methodology

**Trial:** Clinical trials were performed on diabetes patients for 3-days continuously. The sample size of the study was n=55 patients. Medicines were eliminated from the first day of the trial. Thereafter, following 3-days, the participants were kept on a prescribed plant-based diet. Both fasting and post-prandial readings were measured each day along with the weights of the participants. The subjects with varying diabetes history, age groups, type of diabetes, insulin dependency and demographic profiles were part of the trial.

### Findings of the Study

**Results:** The study reported controlled blood glucose levels for 84% of patients and partially-controlled levels for 16% of patients. Those with controlled levels could attain a healthy blood glucose range without medicines and or insulin, **along** with the prescribed diet in 3-days. Those with partially controlled levels could attain a healthy blood glucose range with less than 50% of insulin than prescribed earlier.

Among diabetes type-2 patients the study reported 100% results with all the patients maintaining a healthy blood glucose level. While among diabetes type-1 patients, 57% reported controlled blood glucose levels through the diet and zero-medications. Whereas, 43% reported healthy blood sugar levels through the diet and insulin reduction. In addition of the insulin-dependent group, 59% could completely drop their insulin requirements and 41% could reduce the requirement to at least 50%.

The weight reduction for 55 patients in 3-days was reported as 1.14kgs of average weight loss per individual. Also, the patients had symptomatic relief from general fatigue and weaknesses. The plant-based diet proved to be beneficial with regards to energy and nutritional fulfillments.

### Future Scope

**Conclusion:** Diabetes treatment has both health and economic burden on society. With reference to the present research, a new approach for the treatment of this considered life-style metabolic disability can be shaped. The plant-based diet has been found effective to cure and control diabetes, eliminating the medicine or insulin requirements. Further research on the subject matter can present a medicine-free-food-science based treatment for the disease. At the same time, this unique treatment approach will eliminate the risks of medicine side-effects. On the basis of this research, diabetes education can be developed for better understanding of the disease and better living for the diseased population.

**Keywords:** Plant-Based Diet, Diabetes, Diabetes Type I And Type II, Insulin Dependent, Blood Glucose Levels

## Introduction

In the latest Clinical Practice Recommendations provided by the American Diabetes Association-ADA, Medical Nutrition Therapy is highly recommended for Diabetes care. However, they could not define a dietary pattern or establish a specific nutrition therapy for delay or prevention of the risk of diabetes. They further suggested increased intake of whole grains, fruits, vegetables and legumes, reducing refined and sugary foods (American Diabetes Association, 2015) [1]. In addition, intensive life-style changes were reported to be 58% effective after 3 years by the ADA (American Diabetes Association, 2015).


The Medical Nutrition Therapy for diabetes paved its way through many randomized trials, meta-analysis and observational studies [2]. Improved glycemic and metabolic control were evident with reduction in A1c and blood glucose levels in diabetes patients. The results reported favored both type 1 and 2 population and worked irrespective of disease duration [2]. Moreover, it was also established that life-style interventions work better than metformin in reducing the incidence of type 2 diabetes [3]. The research in hand presents a defined model of managing diabetes with plant-based diet protocol eliminating the need for medications. The study will provide a strong foundation with extensive future scope for research due to its practical implications.

## Review of Literature

The correlation between life-style orientations and diseases has been profound. The present research is rooted within the vast literature present in support of association between diabetes and diet. Some of the related studies have been listed below.

## Glycemic Control and Diabetes

In the study presented by Riccardi et.al on glycemic control in diabetes, they could establish a deep relationship between pre-diabetes and type 1 and 2 diabetes with high glycemic load. Figure 1 shows the effects of high/low glycemic index (GI) on blood glucose levels in type 1 diabetes patients, similar results were seen in type 2 diabetes patients.



**Figure 1:** A 24 weeks analysis on post-prandial blood glucose concentrations obtained in type 1 diabetes patients with low GI-high fiber diet or high GI-low fiber diet (n=63).

**Source:** Riccardi, G., Rivellese, A. A., & Giacco, R. (2008). Role of glycemic index and glycemic load in the healthy state,

in prediabetes, and in diabetes. American Journal of Clinical Nutrition. Available at: <http://ajcn.nutrition.org/content/87/1/269S.full.pdf+html>

The glycemic load is explained as glycemic index (GI) of a particular food multiplied by the amount of carbohydrate contained in an average portion of the food consumed [4]. The study supported low GI and high fiber-rich diet to manage post-prandial blood glucose levels in pre-diabetes and diabetes [4].

Studies also reported that glycemic index can be used as an effective marker along with fiber-content and nutritional value to classify carbohydrate rich foods and their preferences in routine diet [5]. This presented relevance in case of diabetes control and prevention.

In addition, a comparative study for glycemic index or the quantity of carbohydrates on glycated hemoglobin, C-reactive proteins, lipids and plasma glucose on type 2 diabetes patients gave positive results [6]. The outcomes of the 1-year controlled trial on 162 type 2 patients managed by sole diet gave the mean C-reactive protein being 30% low in low-GI diet in a comparison to high GI diet.

The study reported sustainable reductions in post-prandial glucose level and C-reactive proteins and referred the diet management system to aid in type 2 diabetes management [6].

## Animal Protein and Diabetes

The Singapore Chinese Health Study (SCHS) investigated association between dietary patterns and risk of type 2 diabetes in Chinese men and women in Singapore [7]. The study examined 43,176 individuals aged 47-74 years and diabetes free. The study performed Cox regression for diet pattern scores and risk of type 2 diabetes in individuals [8]. Their dietary patterns showed positive relation between meat-rich foods and risk of type 2 diabetes. Whereas fruits, vegetables and soy-rich foods inversely affected risk of type 2 diabetes [8].

Meat consumption and incidence of type 2 diabetes has been elucidated in a cohort study of 4,366 Dutch participants [9]. This study delivered the effects of processed meat on insulin resistivity and incidence of type 2 diabetes. The heating up of meat leads to the formation of AGEs (Advanced glycation end products) [10,11]. It is expected that the pro-inflammatory properties in AGEs may attribute towards the induced risk of type 2 diabetes [12]. Moreover, presence of saturated fatty acids in meat can even contribute to the risk of type 2 diabetes [13].


## Cow's Milk Consumption and Diabetes

Campbell's China Study 2005, a guide to nutrition and health reported that milk protein casein is not fit for human consumption. The proteins in cow's milk have been found to be responsible for auto-immune diseases especially type 1 diabetes; mostly in children with genetic susceptibility [14]. This was explained as, may be in most of us; the body's immune cells are unable to distinguish between the protein fragment of cow's milk and the  $\beta$ -cells of the body. Consequently, the immune cells attack the  $\beta$ -cells of the body resulting in diabetes or other autoimmune diseases [14].

An overview of medical literature on early cow's milk exposure and type 1 diabetes reported an increase of risk factors by

approximately 1.5 times [15]. Higher anti-casein antibodies were also observed in children with type 1 diabetes [16].

In a popular study, a linear model was obtained on analyzing age-standardized prevalence of diabetes among children of 0-14 years of age in 12 countries. The countries were Finland, Sweden, Norway, Great Britain, Denmark, United States, New Zealand, Netherlands, Canada, France, Israel, and Japan (See Figure 2) [17].



**Figure 2:** Association of Cow's milk Consumption and incidence of type 1 diabetes in different Countries

**Source:** Campbell, T. C., & Li, T. M. C. (2005). T. colin campbell, phd and thomas. Available at: <http://www.socakajak-klub.si/mma/The+China+Study.pdf/20111116065942/>

Among them, Finland had the highest incidence of insulin-dependent type 1 diabetes, which was 35 times higher than Japan. Finland has the world's highest cow's milk and milk products consumption and subsequently highest prevalence of diabetes [18]. The research concluded that cow's milk may be responsible for development of insulin-dependent type 1 diabetes.

### Plant-based diet and Diabetes

In support of plant-based diet, a cohort study involved 3,704 participants with 653 diabetes patients from European Prospective Investigation [19]. The study examined the association between intake of fruits, vegetables and fruits and vegetables in combination along with their variety and quantity and risk of type 2 diabetes [19]. The research analyzed 11-year incidence of type 2 diabetes, and reported 21% lower risk of diabetes with greater fruits and vegetables intake in diet [19].

A research based on epidemiological and clinical trials found that nuts can improve post-prandial glycemia and reduce the risk of diabetes [20]. Many studies have reported the relation between nut consumption and metabolic syndrome (MetS). Metabolic Syndrome is a group of cardio-metabolic risk factors, which comprise of type 2 diabetes, high fasting plasma glucose, hyperglycemia, hyper-triglycerides, low HDL cholesterol and abdominal obesity [21].

Metabolic syndrome raises the risk of diabetes by 5 times and that of cardiovascular diseases for diabetes population by 2 to 5 times [21]. Nuts have been found to play an important role in adjusting

the components of MetS by influencing inflammation, oxidative stress, and endothelial function. This in process influences the insulin sensitivity and reduces chances of diabetes, hypertension and obesity [22].

Also, three cohort studies could establish a reduced risk of type 2 diabetes by 27%, 20% and 33% respectively by nut intake in regular diet in women in the Nurses' Health Study (NHS, in the NHS and NHS II cohorts) and women in the Shanghai Women's Health Study [23-25].

A significant research published in the Current Atherosclerosis Reports-2010 demonstrated that, the time of cooking is directly proportional to the increase in glycemic index (GI), resulting in lot of burden on the blood sugar making a person more prone to diabetes [26]. In the same research it was proved that the simple whole grain consumption in its natural state helps the diabetic patient to get a more stable and acceptable blood sugar. However, the refining followed by cooking of the grains rapidly shifts the grains from the low GI range to high GI range [26].

### Research Design

The 3-days Residential Treatment Tour was conducted at Zorba, The Buddha, 10 - Tropical Drive, Ghitorni, New Delhi from 29th April to 1st May, 2016 with 55 diabetes patients and 6 Medical Analysts. The Residential Treatment Tour was publicized among masses both online and through seminars. The procedure required form-filling of DAM form-Diet and Medicine information by the patients. By the time of the beginning of the tour, 60 patients could furnish all details along with completion of the desired formalities of the tour. However, 5 of them had to leave because of personal reasons. Therefore, our sample-size was reduced to 55 patients.

The procedure was planned to keep the 55 subjects on 3-days plant-based raw fruits and vegetables diet. The medicines were discontinued at the start of the plan. Meals were provided as per the diet plan with appropriate quantities based on the patient's weight. Regular blood-sugar readings - fasting and post-prandial were taken and individuals' log-sheets were further maintained. The diet plan was all different for 3-days and was especially designed to fulfill the nutritional requirements of the patients.

### Controlled and Partially Controlled Blood Glucose Levels

Patients who could maintain a healthy reading by the end of 3-days of plant-based diet treatment along with eliminating the requirement for medications were considered as subjects with controlled blood glucose levels.

Those who could maintain a healthy blood glucose level with less than 50% of insulin requirement as was compulsory before, were categorized as subjects with partially controlled levels.

### Findings of the Study

The study reported 46 patients with controlled sugar levels and 9 with partially-controlled sugar levels. The valid percentages obtained were 84% and 16% under controlled and partially controlled groups respectively.

### Sugar Levels

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Controlled	46	83.6	83.6	83.6
	Partially Controlled	9	16.4	16.4	100.0
	Total	55	100.0	100.0	

Frequency Table-1

- Frequency Table 1 - Controlled and Partially Controlled Blood Glucose Levels in 55 Diabetes Patients

1. Correlation Table 1 - Sugar levels and Type 1 or 2 Diabetes in 55 Diabetes Patients

### Sugar Levels \* Diabetes Type Cross-tabulation

		Diabetes Type Total		
		Type 1	Type 2	Total
Sugar Levels	Controlled	12	34	46
	Partially Controlled	9	0	9
Total		21	34	55

Correlation Table-1

**Results:** In the trial 21 patients were type 1 diabetic and 34 patients were type 2 diabetic. Among type 1 patients 57% could attain controlled blood glucose readings and 43% attained partially controlled readings. Among type 2 diabetes patients, 100% gave controlled sugar readings through the process.

2. Correlation Table 2 - Sugar levels and Insulin Dependency in 55 Diabetes Patients

### Sugar Levels \* Insulin Dependency Cross-tabulation

		Insulin Dependency		Total
		Insulin-Dependent	Insulin-Independent	
Sugar Levels	Controlled	13	33	46
	Partially Controlled	9	0	9
Total		22	33	55

Correlation Table-2

**Results:** The trial had 40% insulin-dependent cases, of these 59% could completely drop their insulin requirement to zero and 41% could reduce the levels by at least 50% of the earlier requirement.

3. Correlation Table 3 - Sugar levels and Disease Duration in 55 Diabetes Patients


### Sugar Levels \* Disease Duration Cross-tabulation

		Disease Duration				
		Less than 1 year	1 to 5 years	5 to 10 years	Above 10 years	Total
Sugar Levels	Controlled	25	12	4	5	46
	Partially Controlled	7	1	1	0	9
Total		32	13	5	5	55

Correlation Table - 3

**Results:** An important observation is that 100% results were obtained with patients with above 10 years of disease history, as all the 5 subjects maintained controlled sugar levels. For those newly diagnosed or less than 1 year of disease history, 78% could attain controlled blood glucose readings.

Below is Bar Chart-1 of two variables the attained sugar levels and disease duration in 55 Diabetes Patients.



Bar Chart-1

**Interpretation:** Maximum number of patients were with a disease history of <1 year, 78% of these reported controlled readings, following them were patients within 1-5 years of disease history, who gave 92% controlled results, and 80% controlled results for the group with 5-10 years of disease history.


Frequency Table 2 – Weight Reduction in 55 Diabetes Patients

### Weight Reduction

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Less than 1kg	27	49.1	49.1	49.1
	1 to 2kgs	8	14.5	14.5	63.6
	2 to 3kgs	11	20.0	20.0	83.6
	Above 3kgs	9	16.4	16.4	100.0
	Total	55	100.0	100.0	

Frequency Table-2

**Results:** The findings of the study gave 1.14 kilos of average weight loss per individual of total 55 cases. Among these, 9 subjects could reduce more than 3kilos of weight in 3-days of plant-based diet treatment along with good control over blood glucose levels. Almost half of the cases could reduce <1 kilos of weight during the trial.



Pie-Chart

**Interpretation:** Of the total 55 subjects, 16% reduced  $\geq 3$  kilos of weight, following them were 20% of patients with 2-3kgs of weight reduction and 14% could reduce 1-2 kilos of weight during 3-days. Whereas, 49% reported <1 kilo of weight reduction.

**Correlation Table 4 - Sugar levels and Age Groups in 55 Diabetes Patients**


**Sugar Levels \* Patient Age Cross-tabulation**

Sugar Levels		Patient Age						Total
		Less than 20	20-30	30-40	40-50	50-60	Above 60	
Sugar Levels	Controlled	4	1	11	9	9	12	46
	Partially Controlled	6	1	1	1	0	0	9
Total		10	2	12	10	9	12	55

Correlation Table - 4

**Results:** Among patients with different age-groups, all of those  $\geq 50$  years of age could attain controlled sugar levels. Those below 20 years of age showed 40% controlled and 60% partially controlled sugar levels. This could be related to the little difficulty faced by young children to consume raw-food in those 3-days and report effective results.

Below is the bar-chart plot between two variables-attained sugar levels and weight reduced in 55 diabetes patients.



Bar Chart-2

**Interpretation:** The Bar Chart-2 clearly shows the maximum subjects  $\geq 60$  years of age under controlled group following them are the subjects in 30-40 age group, the 40-50 and 50-60 age groups had equally effective outcomes.

**Correlation Table 5 - Sugar levels and Gender in 55 Diabetes Patients**

**Sugar Levels \* Patient Gender Cross-tabulation**

Sugar Levels		Patient Gender		
		Male	Female	Total
Sugar Levels	Controlled	34	12	46
	Partially Controlled	7	2	9
Total		41	14	55

Correlation Table - 5

**Results:** Of 55 cases, 41 were males and 14 were females. Among them, 83% of males and 86% females could attain controlled blood glucose readings by the end of 3-days diet treatment. The outcomes clearly support the diet protocol to be equally effective in both the genders.

## Conclusion

There had been extensive research on changes in life-style and diet to cure diabetes, but rare could establish a practical approach. Also, most research work is based on one particular type of foods or parameter in terms of diabetes or cardiovascular diseases. Besides this, it is important to relate the nutritional fulfillment through diet in terms of healthy carbohydrates, proteins, fats, vitamins, minerals and anti-oxidants. Furthermore, consideration of body's metabolism through the functional and metabolic pathways can only provide the actual effect of the food in the body.

The plant-based diet protocol has the similar design and works to aid the effective mechanisms in body. Eliminating the toxic components in food, the diet covers all the nutritional requirements.

The plant-based diet in the form of raw fruits and vegetables has the ability to reduce blood glucose levels both fasting and post-prandial. The diet is suitable for diabetic individuals. The diet has shown effect in case of both type 1 and type 2 diabetes patients. The diet process could reduce the insulin dependency for most of the patients by at least 50%. Disease duration was not a hindrance, as similar effects were seen in subjects with above 10 years of disease history and with newly diagnosed diabetes patients.

The diet proved to be beneficial irrespective of variable age groups and demographic profiles. Besides, weight reduction by plant-based diet with quality nutrition makes the diet suitable for diseases associated with obesity or high fatty acids and related metabolic and cardiovascular diseases.

### Post Tour Follow-ups

The necessary follow-ups were practiced post 3-days of Residential Tour. Most of the patients who maintained the diet protocol in their routine gave affirmative response. For most of them their medications completely dropped. Few of them could even maintain a healthy life-style with no-medicines even for common fevers. The remaining, who were still on medicines have been reducing them slowly with the diet-protocol. Follow-ups and advices are still carried over when required.

### Reference - Blood Glucose Upper Limit

In the above report the threshold or diagnostic criteria for blood glucose levels in 55 diabetes patients was taken to be 250mg/dl post-prandial sugar level. This cut-off limit has been established in the book Last-Days of Diabetes [27].

### References

- American Diabetes Association (2015) Standards of Medical Care in Diabetes - 2015 Abridged for Primary Care Providers. *Clinical Diabetes journal* 33: 97-113.
- Pastors J G, Franz M J, Warshaw H, Daly A, Arnold M S (2003) How effective is medical nutrition therapy in diabetes care? *J Am Diet Assoc* 103: 827-831.
- Knowler W C, Barrett-Connor E, Fowler SE, Hamman R F, Lachin J M, et al. (2002) Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin. *The New England Journal of Medicine* 346: 393-403.
- Riccardi G, Rivellese A A, Giacco R (2008) Role of glycemic index and glycemic load in the healthy state, in prediabetes, and in diabetes. *American Journal of Clinical Nutrition* 87: 269-274.
- Mann J I, De Leeuw I, Hermansen K, Karamanos B, Karlström B, et al. (2004) Evidence-based nutritional approaches to the treatment and prevention of diabetes mellitus. *Nutrition, Metabolism, and Cardiovascular Diseases: NMCD* 14: 373-394.
- Wolever T M S, Gibbs A L, Mehling C, Chiasson J L, Connelly P W, et al. (2008) The Canadian Trial of Carbohydrates in Diabetes (CCD), a 1-y controlled trial of low-glycemic-index dietary carbohydrate in type 2 diabetes: No effect on glycated hemoglobin but reduction in C-reactive protein. *American Journal of Clinical Nutrition* 87: 114-125.
- Hankin J H, Stram D O, Arakawa K, Park S, Low S H, et al. (2001) Singapore Chinese Health Study: development, validation, and calibration of the quantitative food frequency questionnaire. *Nutrition and Cancer* 39: 187-195.
- Odegaard A O, Koh W P, Butler L M, Duval S, Gross M D, et al. (2011) Dietary patterns and incident type 2 diabetes in Chinese men and women: the Singapore Chinese Health Study. *Diabetes Care* 34: 880-885.
- Van Woudenberg G J, Kuijsten A, Tigcheler B, Sijbrands E J G, Van Rooij F J A, et al. (2012). Meat consumption and its association with C-reactive protein and incident type 2 diabetes: The Rotterdam study. *Diabetes Care* 35: 1499-1505.
- Hofmann S M, Dong H J, Li Z, Cai W, Altomonte J, et al. (2002) Improved insulin sensitivity is associated with restricted intake of dietary glycoxidation products in the db/db mouse. *American Diabetes Association* 51: 2082-2089.
- Sandu O, Song K, Cai W, Zheng F, Uribarri J, et al. (2005) Insulin resistance and type 2 diabetes in high-fat-fed mice are linked to high glycotxin intake. *Diabetes* 54: 2314-2319.
- Uribarri J, Cai W, Sandu O, Peppas M, Goldberg T, Vlassara H (2005) Diet-derived advanced glycation end products are major contributors to the body's AGE pool and induce inflammation in healthy subjects. *Annals of the New York Academy of Sciences* 1043: 461-466.
- Feskens E J M, Kromhout D (1990) Habitual dietary intake and glucose tolerance in euglycaemic men: The Zutphen study. *International Journal of Epidemiology* 19: 953-959.
- Campbell T C, Li T M C (2005) T. Colin Campbell, Thomas Campbell. <http://www.socakajak-klub.si/mma/The+China+Study.pdf/20111116065942/>.
- Gerstein H C (1994) Cow's milk exposure and type 1 diabetes mellitus: a critical overview of the clinical literature. *Diabetes Care*. 17: 13-19.
- Savilahti E, Akerblom H K, Tainio V M, Koskimies S (1988) Children with newly diagnosed insulin dependent diabetes mellitus have increased levels of cow's milk antibodies. *Diabetes Research* 7:137-140.
- Dahl-Jorgensen K, Joner G H K (1991) Relationship Between Cows' Milk Consumption and Incidence of IDDM in Childhood. *Diabetes Care* 14: 1081-1083.
- LaPorte R, Tajima N, Akerblom H, Berlin N, Brosseau J, et al. (1985) Geographic differences in the risk of insulin-dependent diabetes mellitus: the importance of registries. *Diabetes Care* 8: 101-107.
- Cooper A J, Sharp S J, Lentjes M A, Luben R N, Khaw K T, et al. (2012). A prospective study of the association between quantity and variety of fruit and vegetable intake and incident type 2 diabetes. *Diabetes Care*. 35: 1293-1300.
- Salas-Salvado J, Guasch-Ferre M, Bullo M, Sabate J (2014) Nuts in the prevention and treatment of metabolic syndrome. *American Journal of Clinical Nutrition* 100: 399-407.
- KG Alberti, RH Eckel, SM Grundy, PZ Zimmet, JI Cleeman, et al. (2009) Harmonizing the metabolic syndrome. *Circulation* 120: 1640-1645
- Kendall C W C, Josse A R, Esfahani A, Jenkins D J A (2010) Nuts, metabolic syndrome and diabetes. *The British Journal of Nutrition* 104: 465-473.
- Jiang R, Manson J E, Stampfer M J, Liu S, Willett W C, et al. (2014) Nut and Peanut Butter Consumption and Risk of Type 2 Diabetes in Women. *Journal of The American Medical Association* 288: 2554-2560.
- Pan A, Sun Q, Manson J E, Willett W C, Hu F B (2013) Walnut Consumption Is Associated with Lower Risk of Type 2 Diabetes in Women. *Journal of Nutrition* 43:512-518.
- Villegas R, Gao Y T, Yang G, Li H L, Elasy T A, et al. (2008)

- 
- Legume and soy food intake and the incidence of type 2 diabetes in the Shanghai Women's Health Study. *Am J Clin Nutr* 87:162-167.
26. Murray, P, Chune G W, Raghavan V A (2010). Legacy effects from DCCT and UKPDS: What they mean and implications for future diabetes trials. *Current Atherosclerosis Reports*. Available at: <http://link.springer.com/article/10.1007/s11883-010-0128-1>
27. Chowdhury B R (2016) For Doctors & Care Givers. <https://www.biswaroop.com/9312286540.pdf>.

*Citation:* Biswaroop Roy Chowdhury (2017). *Diabetes Reversal by Plant-Based Diet*. *Journal of Medical & Clinical Research* 2(5):1-7.

*Copyright: ©2017: Biswaroop Roy Chowdhury. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.*