

Endoscopic Findings Related To Helicobacter Pylori

Alexandre Gomes^{1,2*} and Thelma Larocca Skare¹

¹Postgraduate Program in Principles of Surgery, Evangelic Faculty of Paraná/University Evangelic Hospital of Curitiba/Medical Research Institute, Curitiba, PR, Brazil

²Gastrointestinal Endoscopy Service of the 9 of July Hospital, São Paulo, SP, Brazil

Corresponding author

Alexandre Gomes, Postgraduate Program in Principles of Surgery, Evangelic Faculty of Paraná/University Evangelic Hospital of Curitiba/Medical Research Institute, Curitiba, PR, Brazil, E- mail: alex-gomes@uol.com.br

Submitted: 22 Nov 2017; Accepted: 03 Dec 2017; Published: 31 Mar 2018

Abstract

Background: Studies with the latest technologies such as endoscopy with magnification and chromoendoscopy showed that various endoscopic aspects are clearly related to infection by *Helicobacter pylori* (Hp). The description of different patterns of erythema in gastric body under magnification of images revived interest in identifying these patterns by standard endoscopy.

Aim: To validate the morphologic features of gastric mucosa to conventional gastroscopy Hp related gastritis allowing predictability of their diagnosis as well as proper targeting biopsies.

Methods: Prospective study of 339 consecutive patients with the standard videoendoscope image analysis were obtained, recorded and stored in a program database. These images were studied with respect to the presence or absence of HP, diagnosed by rapid urease test and/or by histological analysis. Were studied: a) normal mucosa appearance; b) mucosal nodularity; c) diffuse nonspecific erythema or redness of antrum and body; d) mosaic pattern with focal area of hyperemia; e) erythema in streaks or bands (red streaks); f) raised erosions; g) flat erosions; h) fundic gland polyps. The main exclusion criteria were the use of drugs, Hp pre-treatment and other entities that could affect results.

Results: Applying the exclusion criteria, were included 170 of the 339 patients, of which 52 (30.58%) were positive for Hp and 118 negative. On the positive findings, the most associated with infection were: antralnodularity (26.92%); raised erosion (15.38%) and mosaic pattern in the body (21.15%). On the negative group the normal appearance of the mucosa was 66.94%; red streaks in 9.32%; flat erosions 11.86%; and fundic gland polyps 11.86%. Conclusion: Endoscopic findings are useful in predicting the outcome, localization and targeting of biopsies in gastritis related to *Helicobacter pylori* infection. The most representative form of Hp related gastritis was the nodularity of the antral mucosa. Diffuse redness and mosaic pattern in the body are closely related to Hp infection. Raised erosions are suggestive but not specific to the infection. Normal-appearing forms, red streaks and fundic gland polyposis are related to the negativity of *Helicobacter pylori* infection. Flat erosions were not conclusive of the presence or absence of Hp.

Keywords: Helicobacter pylori, Gastritis, Endoscopy, Gastrointestinal

Introduction

Since the discovery of *Helicobacter pylori* (HP) in 1983, strong evidences have indicated that the infection has an important role in the pathogenesis of chronic gastritis, peptic ulcer and gastric cancer [1]. Gastritis involves cell damage, regenerative and inflammation of the mucosa, with presence of lymphoid follicles. The inflammatory process is initially superficial but, in sequence, affects the entire mucosa, first in the antrum and progressing proximally to the body. Over the years the gastric glands are destroyed, showing epithelial atrophy and intestinal metaplasia areas that favor the appearance of gastric carcinoma [2].

The diagnosis of the infection requires at least two tests in accordance

with the European guidelines [3]. The most used are the rapid urease test and histological analysis [4]. The rapid urease test has a sensitivity of 92% and specificity of 95% [5]. Laine et al found histological sensitivity variation in the identification of Hp bacteria according to the bacterial density in the sample [6]. The H & E staining showed 70% to 98% of sensitivity and specificity of 89% to 98% in the identification of Hp and Giemsa sensitivity of 64% to 96% and specificity of 98% to 100% [6]. Hp is distributed irregularly in gastric mucous epithelial surface and the relatively low density of bacteria in various groups of patients can lead to false-negative results in methods of biopsy [7].

There are few reports in the literature regarding endoscopic patterns of related Hp gastritis using conventional endoscopy. In the initial works in 1995, some authors concluded that it was not possible to establish this diagnosis based on only endoscopy [8,9]. However,

newer technologies such as magnification and chromo endoscopy showed that there endoscopic aspects that are clearly associated with Hp infection, while others relate to uninfected or eradication [10-12]. Yagi et al. described the characteristics of endoscopic findings with magnification in the gastric body with normal appearance and negative Hp: enanthema in tiny streaks or spots in "pinhole" aspect, which correspond to sub-epithelial capillaries and venules networks called RAC (regular arrangement of collecting venules) [13,14]. Anagnostopoulos et al. demonstrated that the mosaic pattern in the gastric body is more related to infection by Hp, as also the enanthema in association with enlarged folds, edema and exudate, indicating intense active inflammatory process [15]. Red streaks and the appearance of tiny red spots corresponding to subepithelial venulescoletantes (normal condition) are associated with the absence of infection by Hp. These authors suggest that, using this technique to perform the biopsy pathology is not required [16].

However, magnification and chromoendoscopy it is not available in most diagnostic centers, and also demand more time for execution and training and does not seem to be practical in daily routine examinations. If specific patterns of Hp-related gastritis can be identified using conventional endoscopy, these patterns could be applied to predict and select patients and biopsies could be targeted to suspected areas of Hp. This study aims to verify the validity of the recognition of morphological patterns of gastritis associated with Hp using conventional endoscopy, which would be helpful to favor the targeting of biopsies for the most affected areas.

Methods

This is an observational cross-sectional study approved by the ethics committee in local research. In this study was analyzed, prospectively, the endoscopic findings of 339 consecutive patients from May 27 2015 until July 10, 2015 in Endoclinic, SP, Brazil. Free and informed consent was obtained from all participants.

The included underwent endoscopy unit with Fujinon 4400 or Pentax EPM 3500 and all tests performed by a single professional. Sedation was carried out with 25 to 50 ug fentanylolamina and midazolam 2-5 mg [17]. Images were recorded and saved in the database (OCRAM® system, SP) being obtained of 12-20 images per patient in all cases. These were selected for this review 6-8 images. Endoscopic aspects that were evaluated were: normal mucosa appearance, nonspecific diffuse erythema of antrum and body, erythema in streaks or bands (red streaks), mosaic mucosal pattern in the gastric body, flat erosions (minor surface defects 5 mm and flat edges), raised erosions, nodularity of the mucosa and fundic gland polyps (Figures 1 & 2).




Figure 1: Endoscopic findings related to the positive HP: A) antral nodularity; B) raised erosions; C and D) gastric body with spotty

redness or mosaic mucosal pattern; E and F) diffuse erythema with edema of gastric folds and thin layer of exudate.




Figure 2: Endoscopic findings related to negative HP: A) normal appearance of the gastric body with regular arrangement of collecting venules; B) red streaks; C) red streaks with flat erosions; D) flat erosions in the antrum; E) fundic gland polyps; F) extensive mucosal atrophy.

The diagnosis of *H. pylori* infection was done by the urease tests (UretestRenylab®, MG) performed with at least two fragments from the antrum and two from the body. The positive histological fragments were subjected to histological examination by hematoxylin and eosin (HE) staining and Giemsa stain to identify the HP, made by a pathologist who was blinded to the other results.

The following exclusion criteria were: patients with anemia, liver cirrhosis, gastric cancer, gastrectomy, renal failure, congestive heart failure, recent use of anti-inflammatory drugs, aspirin, antithrombotics, use of proton pump inhibitors or H2-receptor antagonists in past two months, prior history of eradication of HP; extensive gastric mucosal atrophy.

Applied exclusion criteria were eliminated 169 patients, leaving 170, of which 52/170 (30.5%) were positive and 118/170 (69.4%) Hp negative.

The data were studied in frequency tables and contingency being used the Fisher test and chi-square association for nominal data and Mann-Whitney and unpaired t test for association of numerical data. The significance used was 5% ($p = 0.05$). The calculations were made with the Graph Pad Prism version 5.0 software.

Results

The positive Hp endoscopic findings are shown in Table 1 where it can be seen that the most frequent finding was diffuse redness.

Table 1: Endoscopy findings in 52 patients positive for *Helicobacter pylori* infection.

Findings	n=52	%
Normal mucosa appearance	8	15,38
Antral nodularity	14	26,92
Mosaic pattern in the body	11	21,15

Diffuse redness of antrum and body	30	57,69
Red streaks	0	0
Flat erosions	5	9,61
Raised erosions	8	15,38
Fundic gland polyps	0	0

In Table 2 are the frequency of the findings in the negative group.

Table 2: Endoscopy findings in 118 patients negative for Helicobacter pylori infection.

Findings	n=118	%
Normal mucosa appearance	79	66,94
Antral nodularity	0	0
Mosaic pattern in the body	3	2,54
Diffuse redness of antrum and body	23	21,18
Red streaks	14	11,86
Flat erosions	0	0
Raised erosions	11	9,32
Fundic gland polyps	14	11,86

In about 8/52 (15.38%) infected patients, endoscopic examination was normal while normality was present in 79/118 (66.94%) of the uninfected. Comparing the endoscopic findings in Table 3, it is possible to note that patients with Hp infection have more antral nodularity, mosaic pattern in the gastric body and redness of antrum and body.

Table 3: Comparison of endoscopic findings in HP positive and negative individuals.

Variable	HP positive n=52	HP negative n=118	p	OR	CI
Age years	15- 80 Average of 41.15 ± 14.84	9,0-72,0 Average of 41.14±15.45	0,99 (*)		
Genre	32 women / 26 men	72 mulheres/ 46 men	0,45 (§)		
Normal appearance	8 (15,38%)	79 (66,94%)	<0,0001 (§§)	0,08	0,03-0,20
Nodularity	14 (26,92%)	0	<0,0001 (§§)	OR=89,26	95% CI= 5.19 -153
Mosaic pattern	11 (21,15%)	3 (2,54%)	0,0002 (§§)	OR=10,28	95% CI= 2,73- 38,7
Diffuse redness	30 (57,69 %)	25 (21,18 %)	<0,0001 (§)	OR=5,07	95% CI=2,50- 10,27
Flat erosions	5 (9,61%)	14 (11,86%)	0,33 (§)		
Raise erosions	8 (15,38%)	0	0,008 (§§)	OR=21.9	95% CI=1,16-416
Red streaks	0	11 (9,32%)	0,01 (§§)	OR=0,089	95% CI=0,005-1,54
Polyps	0	14 (11,86%)	0,005 (§§)	OR=0.06	95%CI=0,004 -1,17

(*) = unpaired t test; (§) = chi-square test; (§§) = Fisher test

Discussion

In the present study we sought to identify endoscopy findings related to Hp infected or uninfected gastric mucosa. The selected endoscopic findings for this research have clear association with Hp related gastritis and have been described in previous papers. Edema plies, with or without exudate, diffuse or patchy erythema are regarded as mucosal inflammatory process and are good indicators of the presence of HP [18].

Atrophic gastritis areas were avoided for biopsies in this study because they are hostile regions of the mucosa to the colonization of HP, causing false-negative test results and are present in older patients with longtime HP chronic infection [19].

Magnifying studies have shown that the numerous tiny lines or red dots in the gastric body, seen with conventional endoscopy, were regular arrangement of collecting venules (RAC), characteristic finding in normal stomach without infection by HP with 100% sensitivity and 90% specificity [14].

In the current study the following imaging findings showed a positive association with Hp: antral nodularity, mosaic pattern in the body, diffuse redness and raised erosion.

According to the literature, the antral nodularity is significantly associated with chronic active gastritis and follicular gastritis and showed high specificity (98.5%) and high positive predictive value (91.7%) but low sensitivity (32%) for the diagnosis of HP [8,20]. In this research those images shown to be 89.2 times more frequent in infected individuals (OR = 89.26 and 95% CI = 5.19 -153) demonstrating that this finding is valuable in the diagnosis of Hp infection and the endoscopic finding that best showed this association.

The raised erosions are mucosal elevations on gastric folds of the antrum and distal body containing fibrin exudation and sometimes hematin. Denote chronic inflammation and besides being frequent in patients with Hp, appear also in individuals with chronic use of anti-inflammatory drugs [21]. In this study the findings were nearly 22 times more common in infected patients and none in the negative group (p = 0.0081, OR = 21, 99; 95% CI = 1.16 to 416.6).

The flat erosions are mucosal continuity solutions, associated with erythema, fibrin and sometimes hematin. In general it is smaller than 5 mm in diameter and less than 1 mm depth [21]. These findings were not useful for the diagnosis in question. They represented 9.61% of Hp positive group and 11.86% of HP negative group, with p = 0.33.

The redness of the mucosa was the most common finding. For this study, this kind of finding was divided into diffuse redness in antrum and body, red strikes (according to the literature is found most negative Hp cases) and mosaic pattern (more related in the Hp positive cases) [18]. In this study diffuse redness was found in 30 patients in positive group Hp (57.69%) and in negative group 25 (21.18%), being 5.7 times more common in infected individuals (p <0.0001, OR = 5.07, 95% CI 2.50 to 10.27). The mosaic pattern was found in 11 patients in positive group (21.15%) and in only three patients (2.54%) in the negative group (p = 0.0002; OR = 10.28; 95% CI 2.73 to 38.7 in).

Red streaks findings have shown negative association with infection, but this association was not confirmed by analyzing the confidence

interval obtained. On the negative Hp group were found 11 patients with this aspect (9.32%) and none in Hp positive group ($p = 0.01$; OR = 0.089 95% CI = 0.005 to 1.54).

Fundic gland polyps, according to literature are associated only with uninfected cases. In this work all detected polyps are fundic gland and in all cases Hp was negative ($n = 14$; 11.86%), but this association was not confirmed by analyzing the confidence interval obtained. No hyperplastic polyp (associated with the presence of HP and with congestive gastropathy) was found and also no adenomatous polyp (associated with intestinal metaplasia) [22,23]. 'Polyps fundic gland, red streaks and normal mucosal appearance correlate with the negativity of Hp infection as other studies [19,22,23].

With the selective collection of fragments for histological study in the supposedly positive cases it avoids the routine submission for pathology in cases of morphological patterns not related to infection and with HP negative urease test.

Conclusion

Endoscopic findings are useful in the predictability of the location and direction of biopsies in HP's research. The collection of mucosa fragments in the supposedly positive or negative cases and confirmed by rapid urease test avoids sending routinely for histology in cases of specific morphological patterns.

In the present study, the most representative form of HP related gastritis was the nodularity of the antral mucosa. The raised erosion and mosaic-like appearance in the body were suggestive but not specific to the infection. Fundic gland polyps, red streaks and normal mucosal appearance correlate with the negativity of HP infection. The other forms, diffuse nonspecific erythema or redness of antrum and body and flat erosions, were not conclusive of the presence or absence of HP [24-35].

With the selective collection of fragments, the supposedly positive cases with compatible endoscopic images and urease positive test, avoids the routine sending of material for histology. Similarly, the cases of morphological patterns related to non-infection and urease test negative, the material may be not sent for histological study only in order to confirm the absence of bacteria.

References

1. Komoto K, Haruma K, Kamada T, Tanaka S, Yoshihara M, et al. (1998) Helicobacter pylori infection and gastric neoplasia: correlations with histological gastritis and tumor histology. *Am J Gastroenterol* 93: 1271-1276.
2. Mihara M, Haruma K, Kamada T, Komoto K, Yoshihara M, et al. (1999) The role of endoscopic findings for the diagnosis of Helicobacter pylori infection: evaluation in a country with high prevalence of atrophic gastritis. *Helicobacter* 4: 40-48.
3. Malfertheiner P, Mégraud F, O'Morain C, Hungin AP, Jones R, et al. (2002) Current concepts in the management of Helicobacter pylori infection-the Maastricht 2-2000 Consensus Report. *Aliment Pharmacol Ther* 16: 167-180.
4. Morais M, Macedo EP, Silva Jr MR, Rohr MRS, Ferraz MLG, et al. (1997) Comparação entre testes invasivos para o diagnóstico da infecção pelo Helicobacter pylori. *Arq Gastroenterol* 34: 207
5. Ornellas L C, Cury MS, Lima VM, e Ferrari AP (2000) Avaliação do teste rápido da urease conservado em geladeira. *Arq. Gastroenterol* 37: 155-157.

6. Laine L, Lewin DN, Naritoku W, Cohen H (1997) Prospective comparison of H&E, Giemsa, and Genta stains for the diagnosis of Helicobacter pylori. *Gastrointestinal Endoscopy* 45: 463-467.
7. Khulusi S, Mendall MA, Patal P, Levy J, Badve S, et al. (1995) Helicobacter pylori infection density and gastric inflammation in duodenal ulcer and non-ulcer subjects. *Gut* 37: 319-324.
8. Bah A, Saraga E, Armstrong D, Vouillamoz D, Dorta G, et al. (1995) Endoscopic features of Helicobacter pylori-related gastritis. *Endoscopy* 27: 593-596.
9. Redéen S, Petersson F, Jönsson KA, Borch K (2003) Relationship of gastroscopic features to histological findings in gastritis and Helicobacter pylori infection in a general population sample. *Endoscopy* 35: 946-950.
10. Gonen C, Simsek I, Sarioglu S, Akpınar H (2009) Comparison of high resolution magnifying endoscopy and standard videoendoscopy for the diagnosis of Helicobacter pylori gastritis in routine clinical practice: a prospective study. *Helicobacter* 14: 12-21.
11. Assirati FS, Hashimoto CL, Dib RA, Fontes LHS, Navarro-Rodriguez T (2014) High definition endoscopy and "narrow band imaging" in the diagnosis of gastroesophageal reflux disease. *Arq Bras Cir Dig* 27: 59-65
12. Ratin ACF, Orso IRB (2015) Minimal endoscopic changes in non-erosive reflux disease. *Arq Bras Cir Dig* 28: 20-23.
13. Yagi K, Nakamura A, Sekine A (2002) Comparison between magnifying endoscopy and histological, culture and urease test findings from the gastric mucosa of the corpus. *Endoscopy* 34: 376-381.
14. Yagi K, Nakamura A, Sekine A (2002) Characteristic endoscopic and magnified endoscopic findings in the normal stomach without Helicobacter pylori infection. *J Gastroenterol Hepatol* 17: 39-45.
15. Anagnostopoulos GK, Yao K, Kaye P, Fogden E, Fortun P, et al. (2007) High-resolution magnification endoscopy can reliably identify normal gastric mucosa, Helicobacter pylori associated gastritis, and gastric atrophy. *Endoscopy* 39: 202-207.
16. Ji R, Li YQ (2014) Diagnosing Helicobacter pylori infection in vivo by novel endoscopic techniques. *World J Gastroenterol* 20: 9314-9320.
17. Leslie K, Stonell CA (2005) Anesthesia and sedation for gastrointestinal Endoscopy. *Curr Opin Anaesthesiol* 18: 431-436.
18. Watanabe K, Nagata N, Nakashima R, Furuhashi E, Shimbo T, et al. (2013) Predictive findings for Helicobacter pylori uninfected, -infected and -eradicated gastric mucosa: Validation study. *World J Gastroenterol* 19: 4374-4379.
19. Cho JH, Chang YW, Jang JY, Shim JJ, Lee CK, et al. (2013) Close observation of gastric mucosal pattern by standard endoscopy can predict Helicobacter pylori infection status. *J Gastroenterol Hepatol* 28: 279-284.
20. Laine L, Cohen H, Sloane R, Marin-Sorensen M, Weinstein WM (1995) Interobserver agreement and predictive value of endoscopic findings for H. pylori and gastritis in normal volunteers. *Gastrointest Endosc* 42: 420-423.
21. Appelman HD (1994) Gastritis: terminology, etiology, and clinicopathological correlations: another biased view. *Hum Pathol* 25: 1006-1019.
22. Sakai N, Tatsuta M, Hirasawa R, Iishi H, Baba M, et al. (1998) Prevalence of Helicobacter pylori infection in patients with hamartomatous fundic polyps. *Dig Dis Sci* 43: 766-772.
23. Elhanafi S, Saadi M, Lou W, Mallawaarachchi I, Dwivedi A, et al. (2015) Gastric polyps: Association with Helicobacter

- pylori status and the pathology of the surrounding mucosa, a cross sectional study. *World J Gastrointest Endosc* 7: 995-1002.
24. Kato T, Yagi N, Kamada T, Shimbo T, Watanabe H, et al. (2013) Diagnosis of *Helicobacter pylori* infection in gastric mucosa by endoscopic features: a multicenter prospective study. *Dig Endosc* 25: 508-518.
 25. Dixon MF, Genta RM, Yardley JH, Correa P (1996) Classification and Grading of Gastritis: The Updated Sydney System in the International Workshop on the Histopathology of Gastritis, Houston 1994. *American Journal of Surgical Pathology* 20: 1161-1181.
 26. Loffeld RJ (1999) Diagnostic value of endoscopic signs of gastritis: with special emphasis to nodular antritis. *Neth J Med* 54: 96-100.
 27. Misra SP, Dwivedi M, Misra V, Agarwal SK, Gupta R, et al. (1990) Endoscopic and histologic appearance of the gastric mucosa in patients with portal hypertension. *Gastrointest Endosc* 36: 575-579.
 28. Zerbib F, Vialette G, Cayla R, Rudelli A, Sauvet P, Bechade D, et al. Follicular gastritis in adults. Relations with *Helicobacter pylori*, histological and endoscopic aspects. *Gastroenterol Clin Biol* 17: 529-534.
 29. Watanabe M, Kato J, Inoue I, Yoshimura N, Yoshida T, et al. (2012) Development of gastric cancer in nonatrophic stomach with highly active inflammation identified by serum levels of pepsinogen and *Helicobacter pylori* antibody together with endoscopic rugal hyperplastic gastritis. *Int J Cancer* 131: 2632-2642.
 30. Lopes AI, Vale FF, Oleastro M (2014) *Helicobacter pylori* infection - recent developments in diagnosis *World J Gastroenterol* 20: 9299 -9313.
 31. Yan SL, Wu ST, Chen CH, Hung YH, Yang TH, et al. (2010) Mucosal patterns of *Helicobacter pylori*-related gastritis without atrophy in the gastric corpus using standard endoscopy. *World J Gastroenterol* 16: 496-500
 32. Magalhães, MAB,; Barbosa AJA, Figueiredo JA, Alberti LR, Petroianu A (2015) Effects of different periods of gastric ischaemia in the viability of the tissue of body, fundus and antrum region of rabbit stomach.. *Arq Bras Cir Dig* 28: 167-170.
 33. Henry MACA (2014) Diagnosis and management of gastroesophageal reflux disease. *Arq Bras Cir Dig* 27: 210-215.
 34. Bahú M G, Silveira TR, Maguilnick I, Ulbrich-Kulczynski J (2003) Endoscopic nodular gastritis: an endoscopic indicator of high-grade bacterial colonization and severe gastritis in children with *Helicobacter pylori*. *J Pediatr Gastroenterol Nutr* 36: 217-22.
 35. Calabrese C, Di Febo G, Brandi G, Morselli-Labate AM, Areni A, et al. (1999) Correlation between endoscopic features of gastric antrum, histology and *Helicobacter pylori* infection in adults. *Ital J Gastroenterol Hepatol* 31: 359-365.

Citation: Alexandre Gomes and Thelma Larocca Skare (2018). *Endoscopic Findings Related To Helicobacter Pylori*. *Journal of Medical & Clinical Research* 3(1):1-5.

Copyright: ©2018 Alexandre Gomes. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.