

ISSN: 2577 - 8005

Medical & Clinical Research

Vitamin D, Calcium to Magnesium Ratio, and the Gut Microbiome

Patrick Chambers*

Torrance Memorial Medical Center, Department of Pathology, 3330	[•] Corresponding Author
Lomita Boulevard, Torrance, CA 90505, USA	Patrick Chambers, Torrance Memorial Medical Center, Department of
	Pathology, 3330 Lomita Boulevard, Torrance, CA 90505, USA.

Submitted: 07 Feb 2025; Accepted: 11 Feb 2025; Published: 25 Feb 2025

Citation: Chambers P. (2025). Vitamin D, Calcium to Magnesium Ratio, and the Gut Microbiome. Med Clin Res, 10(2), 01-14.

Abstract

The gut microbiome has been the subject of increasing interest as integral to our health. Few realize that the enormous benefits of vitamin D (VD) and magnesium (Mg) are highly dependent on a healthy gut microbiome. Short chain fatty acids, especially butyrate, reflect not only a healthy gut microbiome but also VD status. Suboptimal VD, Mg, or butyrate translates to some degree of gut dysbiosis and vice versa. Mg dependent secondary bile acids, indoles, and tryptophan, all microbial metabolites and longevity agents, are also discussed. Mg is indispensable to not only the synthesis of the active form of VD but also that of 7-dehydrocholesterol (7-DHC) from acetate. 7-DHC is the substrate for solar conversion to D3. The steadily increasing *Ca:Mg in the Western diet and its ironic impact on parathormone (PTH) is discussed. Gut dysbiosis further complicates this.* Biochemical and physiologic interlinkages are legion and most remain hidden. This limited mini review exposes insight into the tight linkage between 25(OH)D and Ca:Mg, facilitated by the gut microbiome. A model incorporating the physiologically discordant but reinforcing effects on this linkage based on genes, culture, socioeconomic status, and diet that also addresses the seemingly contradictory reports regarding calcium (Ca), Mg, and VD efficacy is proposed. The balance between Ca and Mg is at the heart of the global epidemic of insulin resistance. A reliable method for estimating intracellular Mg++ is also proposed. A primary goal of this paper is to awaken the global public to not only the inadequacy of the RDA for Mg dependent vitamin D but also to that for the RDAs for Mg and Ca. Using generally accepted serum reference ranges, the difference between RBC Mg and plasma Mg, and basic physiology, one can both uncover the serious shortfall involving all three and unequivocally demonstrate the dire health risks linked to an out of range Ca:Mg. Conclusions are observational not experimental.

Keywords: Butyrate, 25(OH)D, Intracrine, Socioeconomic, Kynurenine

1. Introduction

The term VD is often used indiscriminately. VD in this mini review will be used collectively to include its three forms D3 (cholecalciferol), 25(OH)D (storage form), and 1,25(OH)2D (active form). VD, Mg, and the products of a healthy gut microbiome, e.g., short chain fatty acids (SCFAs), secondary bile acids, and indoles, are all longevity agents [1-4]. They are also intertwined both directly and indirectly. VD regulates Mg status as well as that of Ca and phosphate. Many are familiar with the total dependence of VD efficacy, whether of solar or supplemental origin, on adequate Mg. But the vital role of the gut microbiome in potentiating both has only recently been revealed. Gut dysbiosis disrupts the balance of beneficial bacteria and impedes the production of many vital nutrients [5]. Adequacy of the storage form of VD has often been measured relative to parathormone (PTH). But recent research challenges this and raises questions about the impact of Ca:Mg on optimal VD efficacy.

2. Discussion

2.1.Vitamin D and Magnesium

VD and Mg are inextricably linked in a bidirectional manner. Through PTH VD can regulate the intestinal absorption and urinary excretion of Ca and Mg [6]. However, Mg is indispensable to the synthesis of VD. It is generally known that Mg is a required cofactor for every enzymatic step in the conversion of D3 aka cholecalciferol to its active form 1,25(OH)2D, including the binding of D or 25(OH)D to VDBP (vitamin D binding protein). However, Mg is also required for the synthesis and cAMP mediated secretion of PTH from chief cells in the parathyroid gland [7,8]. Low plasma Ca driven PTH synthesis stimulates VD synthesis and VD driven high plasma Ca inhibits PTH synthesis. Even more importantly the synthesis of 7-dehydrocholesterol (7-DHC), the immediate precursor of cholecalciferol, from acetate is dependent on Mg (Figure 1). Acetate is provided by either gut microbes or acetyl CoA, which requires Mg dependent B5 (pantothenate). Vitamins B2 and B3 must be phosphorylated to FAD and NAD respectively to attain active status for the synthesis of 7-DHC. This phosphorylation requires ATP and Mg.

Without sufficient 7-DHC the sun is powerless to create D3. Mg applied in any form enhances skin health. But this is overlooked in favor of VD, despite the fact that solar D3 requires Mg and the active form of VD requires Mg [9]. Both forms play a vital role in protecting the integumentary system, but few articles acknowledge their total dependence on Mg [10].



Figure 1: All enzymes that produce the active form of VD $(1,25(OH)_2D)$ from D3, including binding to the transport protein, synthesis of PTH, and secretion of PTH, are Mg dependent. Many enzymes and cofactors (B2,B3) involved in the synthesis of 7-dehydrocholesterol from acetate are also Mg dependent. 7-DHC=7-dehydrocholesterol, VDBP=vitamin D binding protein

Optimal 25(OH)D is at least 50 ng/mL (120 nmol/L), based on clinical data (Figure 2). The 20 and 30 ng/mL 25(OH)D targets generally recommended are suitable only for rickets and skeletal health, but optimal immune function involves intracrine, autocrine,

and paracrine pathways and storage form levels that exceed those adequate for endocrine (hormonal) needs. Unfortunately 25(OH)D levels lower than 20 ng/mL are considered deficient and those less than 30 ng/mL are only considered insufficient.



Figure 2: Adapted by Robin Whittle May 2022 from two studies [11-13]. The two colored curves represent VD associated risks for surgical site infection and hospital acquired infection (immune functions of VD) from the gastric bypass study [12]. The inadequate D3 supplementation distribution figure is from a Covid-19/VD study [13].

The RDA (Recommended Daily Allowance) of 600–800 IUs D3 per day recommended by the National Academy of Sciences Institute of Medicine (IOM) was proven to be off by an order of magnitude in 2014 [14]. This huge error was confirmed by several Canadian and American university research teams. The IOM's RDA for VD still stands at 600 IUs to 800 IUs (for adults over 70), but their estimated average requirement (EAR) was subsequently raised from 20 ng/mL to 30 ng/mL. EAR + 2SDs = RDA. An increased intake of an order of magnitude, e.g., 8000 IUs of D3 (cholecalciferol), would correlate with a serum level of at least 50 ng/mL [15].

NHANES (National Health and Nutrition Examination Surveys) data between 1988 and 2006 revealed little change in mean serum 25(OH)D at about 25 ng/mL [16]. From 2011 to 2018 median 25(OH)D went from 27 to 27.5ng/mL [17]. Regarding the cations Ca⁺⁺ and Mg⁺⁺ in the relatively healthy (non-hospitalized) with normal renal function without GI issues and no impacting medications, Ca++ is usually about 50% of serum values and Mg+ is usually about 70% of serum values (section 2.2). When midrange values for lab reference limits for Ca and Mg are compared using these percentages, the resulting serum ratio for iCa:iMg aligns well with 2.0 (mmol comparison). Jean Durlach, founder of the International Society for the Development of Research on Magnesium (SDRM) in the early 70's, stated in 1989 that optimal Ca and Mg balance was physiologically 2:1 [18]. His designation of 2.0 was based on research into the physiological roles and interactions of these minerals in the body, not on a questionnaire. Serum Mg is not even offered on a routine chemistry panel. Furthermore, in order to avoid normomagnesemic Mg deficiency aka chronic latent Mg deficiency, the lower limit of the normal range for Mg should be raised from 0.75 to 0.85 mM [2,19-23]. In one study trial participants completed a dietary questionnaire that indicated suboptimal Mg status in 100% of participants. Yet 25% were found to have optimal serum status [24]. Suboptimal Mg status was defined as serum Mg<2.0 mg/dL (0.83 mM). Even increasing the lower limit of normal from 0.75 mM to 0.83 mM does not appear to exclude the 25% with optimal serum Mg judged to be suboptimal by the food frequency questionnaire (FFQ). Furthermore, serum Mg is difficult to sustain with increasing age/ BMI and decreasing food quality.

2.2 Calcium to Magnesium Ratio

NHANES has not determined serum magnesium levels in its participants since 1974 [25]. Since that date the RDA in mg/d has been derived from a FFQ that has been configured to accurately reflect lab values in mmol/L that are clinically relevant. The results from the NHANES provide the raw data for this. The RDA for Mg was 6 mg/kg in developed countries until 1997. Mildred Seelig, coauthor with Andrea Rosanoff of The Magnesium Factor, had previously endorsed an RDA of 6-10 mg/kg/day for young adults and twice that for those under stress. Instead in 1997 the RDA was lowered to 4.5 mg/kg and has not been updated since. Since the RDA for males is 420 mg/d, this recommendation requires that he be 200 lbs (93 kg), the mean weight of an adult male. Even

at this recommended level half fall short. Magnesium intake is inversely associated with obesity risk. According to NHANES data, overweight and obesity have been steadily increasing with nearly 75% of Americans now in this category. Over 40% are obese (BMI > 30). In 1989 Durlach reported his physiological determination that Ca:Mg should target 2.0 based on iCa:iMg. That the ionized forms are more accurate has been reiterated for both [26,27]. He recommended 6 mg/kg. Many articles that touch on the Ca:Mg ratio report their total serum concentrations in mg/ dL, when unbound millimolar concentration would be much more physiologically accurate.

Ca and Mg compete for the calcium sensing receptor (CaSR), and PTH responds to both cations in the same way but not to the same degree [28]. Yet Mg⁺⁺ often opposes Ca⁺⁺, e.g., as a Ca channel blocker. Because serum Ca++ is more responsive to VD and the primary determinant of Mg dependent PTH versus Mg++, an unbalanced intake of these competing cations can create a dilemma that is almost predictable. Furthermore, the lower limit of normal for serum Mg++ is a level insufficient for adequate PTH synthesis, creating a catch 22 [29,30]. A diet induced elevation of Ca:Mg suppresses PTH, depressing Mg absorption and increasing magnesuria. Mg is an "innocent bystander" to this unbalanced (increased Ca:Mg) driven suppression of PTH. Mg dependent PTH synthesis and secretion further compromise the Mg shortfall and facilitate the dietary Ca:Mg imbalance. The Occidental diet is often relatively short Mg versus Ca and the Oriental diet is often relatively short Ca versus Mg. There is a global shortfall of both calcium and magnesium, but the latter is much more pronounced in developed countries. Ca:Mg in the Oriental diet is little more than half that in the Occidental diet, because the Oriental diet contains more magnesium. Poor nutrition and low socioeconomic status in either may be primary determinants of the ultimate ratio [31].

Routine determination of serum ionized magnesium appears to be problematic. Total Mg (bound and unbound) is not even offered on routine chemistry panels. However, by knowing the range for total serum Mg in mg/dL and the range for ionized Mg in mmol/L by ion specific electrode in the relatively healthy, one can show that about 70% of total Mg must be ionized for alignment [32]. Total serum Mg range of 1.8-2.2 mg/dL yields serum Mg⁺⁺ range of .53 - .64 mmol/L. Direct measurement of Mg⁺⁺ by ionic specific electrode reveals the normal range to be .54 - .67 mmol/L [32].

If the following are assumed

1. Intestinal absorption of Mg is 30 to 40% of intake (generally accepted range)

2. Atomic weight of Mg is $24.3 \Rightarrow 24.3 \text{ mg} = 1 \text{ mmole of Mg}^{++}$

3. The reference range for RBC Mg is 3.7-7.0 mg/dL or 1.52-2.88 mmol/L, i.e., mean of 2.2 mmol/L [33]

4. The reference range for plasma Mg⁺⁺ is .54-.67 mmol/L, i.e., mean of .6 mmol/L [32]

5. The reservoir of RBC Mg $^{\!\!+\!\!+}$ is about (2.2/.6=3.67) times serum Mg $^{\!\!+\!\!+}$

6. Plasma volume is about 3 liters and RBC volume is about 2

liters

7. Approximately 70% of serum Mg is unbound

8. An accepted normal range for serum Mg at 1.8-2.2 mg/dL

then, assuming homeostatic equilibrium, one can show that an intake of 400 mg (RDA recommends 310 mg/d for females and 420 mg/d for males) of elemental Mg/d at 35% intestinal absorption will yield a serum Mg⁺⁺ of about .40 mmol/L, i.e., hypomagnesemia. The IOM and the NIH in extrapolating FFQ derived Mg intake (mg/d) to expected plasma Mg (mg/dL) appear to have assumed that RBC Mg is only twice serum Mg⁺⁺ (versus well over 3.0) or that much more than 35% is absorbed. At 40% absorption plasma Mg⁺⁺ increases to .45 mmol/L (hypomagnesemia). Even at 50% absorption plasma Mg⁺⁺ only increases to .56 mmol/L. But it seems that more than 40% absorption would require some assistance from Mg dependent VD. However, Mg intake in those with Mg⁺⁺ at the lower limit of normal (0.54 mmol/L) appears to be allocated first to PTH synthesis and then to VD synthesis [29,30].

The accepted normal range of .75 mmol/L - .95 mmol/L for total serum Mg (.54-.67 mmol/L for unbound) is based on a sampling of the "normal" population, half of which are magnesium deficient. Values between .75 and .85 mmol/L for serum Mg are insufficient, at least in the absence of sufficient Mg dependent VD synthesis. Raising the lower limit to .85 mmol/L enables an Mg⁺⁺ of about .60, facilitating sufficient VD synthesis. Given the differential between RBC magnesium and serum Mg⁺⁺, the RDA for magnesium intake should be at least 570 mg/d. This translates to .56 mmol/L of Mg⁺⁺. To attain .60 mmol/L of Mg⁺⁺ would require an RDA of 610 mg/d. Fear of the laxative effect of Mg⁺⁺ may discourage this. If the following are assumed

1. Intestinal absorption of Ca is about 35% of intake [34]

- 2. Atomic weight of Ca is $40 \Rightarrow 40 \text{ mg} = 1 \text{ mmole of Ca}^{++}$
- 3. plasma volume is 3 liters for one weighing 150 lbs
- 4. The normal range for total serum Ca is about 8.5-10.5 mg/dL
- 5. The normal range for iCa is about 4.5-5.6 mg/dL

6. Approximately 53% of serum Ca is unbound

7. The accepted range for serum Ca⁺⁺ is 4.5-5.6 mg/dL = 1.125-1.4 mmol/L, i.e., a bit higher [35].

then, assuming homeostatic equilibrium, one can show that an intake of 1000 mg (RDA recommends 1000 mg/d for those 19-50 and 1200 for females over 50 and males over 70) of Ca/d at 35% intestinal absorption will yield a serum Ca⁺⁺ of 1.55 mmol/L, which divided by the RDA derived plasma Mg⁺⁺ of .40 yields 3.6 generated by RDA. In summary the intake of Ca and Mg recommended by the IOM and the NIH leads to a physiologic Ca⁺⁺:Mg⁺⁺ of 3.6 much higher than the 2.0 recommended by Durlach. When the midpoints for the established serum ranges for the ionized forms, 0.6 mM for Mg⁺⁺ and 1.26 mM for Ca⁺⁺, the iCa:iMg is 2.1, very much in line with Durlach's physiologic recommendation [32,35].

Regarding the optimal range for Ca:Mg, two studies on Ca and Mg intakes (FFQ) favored an upper limit of 2.8 ratio [36,37]. These studies demonstrated increased risk for colorectal tumors and prostate cancer when Ca:Mg exceeded 2.8, i.e., the ratio was clinically determined and based on actual physiologic iCa:iMg.

Some studies suggest that the upper limit of normal be reduced to 2.6 or even 2.5 based on clinical data [19,38].

Regarding the lower limit, a study on a large Chinese population recommended a median dietary Ca:Mg of 1.7 as a lower limit [31]. This was corroborated by two subsequent Chinese studies that found actual serum reference ranges for Ca:Mg to be between 2.4 and ~3.6 [39,40]. These serum mmol/L levels translate (70% Mg and 53% Ca unbound) to iCa:iMg between 1.7 and 2.5 or mean ~2.1 (Figure 5). Diet is more plant based versus the Western diet.

Therefore, the lower limit of the target range for Ca:Mg intake (mg/d) (determined in the relatively more Ca deficient East) should be 1.7, while the upper limit (determined in the Ca and Mg deficient West) should be less than 2.8 [41]. Yet the mean ratio of Ca:Mg in the US by FFQ over the past 20 years has exceeded 3.1. That derived from the RDAs for Ca and Mg is 3.6, using 1000 mg/d and 400 mg/d RDAs for Ca and Mg respectively.

In summary, based on Chinese studies that correlate intake with serum levels, FFQ determined Ca:Mg appears to more accurately reflect serum iCa:iMg in determining mortality and representing actual clinically relevant iCa:iMg [24]. Several Western studies correlating clinical status with FFQ determined Ca:Mg suggest that the upper limit may be as low as 2.5 [31,39,40]. A rural Eastern diet threatens the lower limit and an urban Western diet threatens the upper limit. Both are short Ca, but the former has more Mg and a lower Ca:Mg. Any meta-analysis that doesn't differentiate is conflicted. The global popularity and growing availability of phosphate laden processed meats and soft drinks undoubtedly are much to blame (Section 2.6).

Regarding supplementation, note that the labels on most Mg containing supplements report the % RDA by serving size, which may not be one tablet. Furthermore, elemental magnesium usually comprises about 10% of the ingested form, and, unless stated otherwise, that's the extent of elemental magnesium in the supplemental tablet.

2.3 Calcium to Magnesium and Vitamin D

Any Ca and Mg discussion must include VD. Serum Ca++ triggers negative feedback inhibition of PTH release that directly impacts VD synthesis and VD mediated Ca and Mg intestinal absorption and renal resorption. PTH and VD status are inversely or negatively correlated and produce a hyperbolic curve (Figure 3). However, might the Ca:Mg impact this relationship, given the almost predictable physiologic dilemma previously discussed? Might the secondary hyperparathyroidism in figure 3 be due to hypocalcemia or a low Ca:Mg? On the other hand, might an elevated Ca:Mg create some degree of secondary hypoparathyroidism (Figure 4) not addressed in figure 3? Might a Mg shortfall, compromising PTH and VD synthesis, obscure a 25(OH)D level of 50 ng/mL obtained primarily through D3 supplementation?



Figure 3: This figure is from Ginde et al. [42]. Any 25(OH)D value < 50 ng/mL implies some degree of secondary hyperparathyroidism.

One relevant study on this [29] reported that when baseline 25(OH) D exceeded 30 ng/mL, Mg supplementation suppressed 25(OH) D. PTH is not mentioned in that study. However, the Ca:Mg ratios for the placebo and target groups were elevated at 3.9 and 3.7 respectively. This suggests that either Mg supplementation when Ca:Mg is elevated may regrettably suppress PTH and with it VD synthesis or that the additional Mg intake (mean 200 mg) may have exceeded bowel tolerance in a significant number of participants. Any Mg supplementation when Ca:Mg is elevated and 25(OH)D levels exceed that sufficient for skeletal health (30 ng/mL) may be counterproductive and may compromise immune health. Perhaps in addressing an elevated Ca:Mg lowering Ca intake should precede Mg supplementation. The resistance by the IOM and NIH to increasing the lower limit of the reference range for Mg from 0.75 to 0.85 mmol/L may be fear of its laxative effect, which facilitates loss of K+ and Mg++. If excess Ca intake is not addressed first, then any increase in Mg intake must still compete for the few available intestinal CaSRs, making the laxative effect hard to avoid.

Other studies support the bell shaped figure 4 conjecture (Figure 5). Lung cancer and CVD are increased whether Ca:Mg is high

[43,44] or low [45]. VD levels are low in both lung cancer and CVD [46,47]. So, the highly touted benefits of VD may be compromised in those with an unbalanced Ca:Mg. However, a low Ca:Mg and insufficient Ca should trigger PTH secretion, but restrictions due to genes, e.g., skin pigmentation, to culture, e.g., Asian umbrellas, female Arab attire, to socioeconomic status, e.g., poor nutrition, and/or to diet, e.g., lactose intolerance, may conspire to deny the intake of Ca or VD or synthesis of D3 dependent VD = depressed 25(OH)D, elevated PTH (see figure 3) => secondary hyperparathyroidism. However, the inverse or curvilinear relationship between PTH and 25(OH)D (left wing in figure 4) may pivot around the optimal zone (see figure 5) with transition to a direct or linear relationship (right wing in figure 4)=> secondary hypoparathyroidism. The vertical axis in figure 4 may also directly reflect gut microbiome quality and optimal health (Figure 5).

So, some normomagnesemics might also be VD deficient with low PTH, i.e., low Mg induced hypoparathyroidism. Any D3 supplementation in the absence of additional Mg would characterize the right wing versus the elevated PTH hyperparathyroid left wing. A low dietary Ca intake, e.g., Chinese diet, upregulates PTH which primarily increases Ca absorption/resorption relative to Mg. But the low intake of Ca should leave more CaSRs available and Mg is somewhat more easily absorbed, slightly increasing PTH synthesis but suppressing VD synthesis, creating a hypoparathyroid low VD scenario. Any supplementary D3 should not only exacerbate this but might also theoretically enable 50 ng/mL 25(OH)D3 levels in the face of a persistent Mg⁺⁺ "normomagnesemic" shortfall. Furthermore, Mg levels decrease with age and with increasing BMI.

Figure 4 represents a hypothetical view of this based on

1. Increasing Ca intake when Ca:Mg is less than 1.7 decreases risk for some cancers [48].

2. Increasing Ca intake when Ca:Mg is greater than 2.8 increases risk for colorectal cancer [36].

3. Increasing Mg intake when Ca:Mg is less than 1.7 increases risk for some cancers [48].

4. Increasing Mg intake when Ca:Mg is greater than 2.8 decreases risk for prostate cancer [37].

5. An elevated Ca to Mg ratio increases risks for some cancers, including lung cancer, and CVD [43,44].

6. A depressed Ca to Mg ratio increases risks for some cancers, including lung cancer, and CVD [44,45].

7. Low Mg in the setting of elevated Ca:Mg translates to low VD. This can be explained physiologically, as Mg is required for the synthesis and secretion of PTH and for the synthesis of VD. Elevated Ca also displaces Mg from CaSRs.

8. Low Ca in the setting of depressed Ca:Mg is physiologically contradictory to a concomitant low VD. However, this may be explained based on discrepant but mutually reinforcing genetic, cultural, socioeconomic, and dietary considerations. These may complicate and compromise clinical correlations and data analysis. 9. Skin pigmentation is directly linked to VD deficiency [49].

10. Socioeconomic status is directly linked to VD deficiency [50,51]. D3 is not in the budget or on the menu.

11. Cultural customs can drive VD deficiency. Most in the Middle East dress modestly and many Asians are averse to solar exposure [52].

12. Diet is largely dependent on culture. The South Asian diet is low in VD rich foods and many Asians are lactose intolerant and avoid dairy products, excellent sources of Ca and VD [53,54].



Ca:Mg e proposed relationships

Figure 4: This hypothetical bell shaped curve delineates the proposed relationships between VD, Ca:Mg, and PTH. Yellow represents the target range and orange the health risk ranges for Ca:Mg, when physiologic considerations and genetic, cultural, socioeconomic, and dietary considerations are separated. The left wing is suboptimal and represents some degree of secondary hyperparathyroidism due to hypocalcemia (Ca<lower limit, Mg</=lower limit), while the right wing is also suboptimal and represents some degree of secondary hyperparathyroidism due to chronic latent Mg deficiency or hypomagnesemia (Ca</=lower limit, Mglower limit). Figure 3 recognizes only the left wing of the bell in figure 4.

Figure 4 illustrates a proposed shift in VD supplementation from one viewed through PTH to one viewed through Ca:Mg. The balance between Ca and Mg is increasingly recognized as critical to attaining optimal levels of 25(OH)D. The left wing represents some degree of secondary hyperparathyroidism due to hypocalcemia, while the right wing represents some degree of secondary hypoparathyroidism due to chronic latent Mg deficiency or hypomagnesemia. Insufficient Mg translates to insufficient Mg dependent PTH synthesis, insufficient PTH induced Ca absorption/resorption, and insufficient Mg dependent VD synthesis => depressed 25(OH)D and PTH. Increasing Mg intake should elevate both 25(OH)D and PTH. But this doesn't happen, when 25(OH)D exceeds 30 ng/mL and Ca:Mg exceeds 3.7 by FFQ (mg/d) (29).



Figure 5: This figure is from Rosanoff et al. [41]. Disease risk, an inverse proxy for VD status, increases as the Ca:Mg increases or decreases. However, increasing Mg with Ca:Mg greater than 2.8 and 25(OH)D greater than 30 ng/mL may not reduce risk, unless excess Ca intake is addressed first [29].

Many Americans may be susceptible to health risks linked to hypoparathyroidism and low Mg dependent PTH that may compromise the efficacy of VD (Figures 4 and 5), especially if 25(OH)D exceeds 30 ng/mL [29].

According to the NHANES from August 2021-August 2023, the Ca:Mg for both genders over 20 years old was 3.01 (diet, no supplements) [55]. According to NHANES 2001-2006 the mean supplementation for adults of both genders was 141 mg/d Ca and 39 mg/d Mg [56].

Extrapolating this supplementation data from NHANES 20001-2006 increases the Ca:Mg to 3.1 by FFQ (mg/d). This is the lowest ratio over the previous two decades. Ca:Mg ratios over this time are much worse for those 6-19 years of age, usually exceeding 4.0 [54].

VD deficiency may be at the root of many health issues exacerbated by a Ca:Mg ratio outside the 1.7-2.8 range [41] (Figure 4). The failure to normalize for Ca:Mg in any study on the efficacy of VD may compromise any conclusions and may sell short the value of this micronutrient extraordinary (Figure 4). The Covid-19 epidemic demonstrated the importance of this ratio, as low Mg intake was tightly linked to Covid severity [57]. A pre-Covid study on high dose D3 for critically ill patients reported no benefit [58]. Ca:Mg status was not reported and magnesium was not mentioned.

Might D3 therapy exacerbate an elevated Ca:Mg?

The Covid-19 epidemic is dwarfed by the ongoing and escalating global pandemic of insulin resistance, as is the prevalence of prediabetes [59]. Reports on insulin resistance vary from 16% to 47% of the population [60]. There is also a silent epidemic of diabetes in low and middle income countries involving about 40% of the population Magnesium deficiency aggravates insulin resistance and Mg supplementation enhances insulin sensitivity and decreases insulin resistance [61-63]. Mg++ is required for activation of the insulin receptor, cAMP mediated release of insulin, and metabolism of glucose [64]. Insulin resistance leads to pre-diabetes and ultimately T2DM.

2.4 Vitamin D and the Gut Microbiome

Although the active form of VD mediates its endocrine, intracrine, autocrine, and paracrine effects, D3 contributes additional benefits. D3 inhibits Candida hyphal morphogenesis [65,66]. The emerging role of the gut microbiome in health and disease and the impact of Candida overgrowth in gut dysbiosis highlights the value of D3. Candida overgrowth can be both cause and effect of gut dysbiosis [67]. In addition the active form 1,25(OH)2D, but not its storage form 25(OH)D, most frequently requested lab analyte, is linked to an abundance of butyrate producing bacteria, frequently linked to gut health [68]. In fact the active form of VD correlated even more tightly with microbiome diversity. These reported results

were adjusted for solar exposure. The active form of VD increases the Bacteroidetes/Firmicutes ratio, especially Akkermansia and Faecalibacterium of the Bacteroidetes phylum, both prominent butyrate producers, and increases microbial diversity [69,70]. VD deficiency and suboptimal gut microbiome are associated with cancer, autoimmune disease, inflammatory bowel disease, cystic fibrosis, multiple sclerosis, diabetes, and depression [71-78]. Butyrate has an ameliorative effect on dementia, cancer prevention/treatment, especially colorectal cancer, and obesity [79-82]. The SCFAs acetate, propionate and butyrate are produced from the microbial fermentation of indigestible carbohydrates and are the biomarkers of a healthy gut microbiome [83]. Acetate and propionate producing bacteria can cross feed butyrate-producing bacteria [84].

But improvements in the gut microbiome via increasing D3 supplementation in those already deficient/insufficient are not limited to the symptomatic. Increased serum 25(OH)D was associated with increased beneficial bacteria and decreased pathogenic bacteria [85]. D3 is both therapeutic and prophylactic and appears to possess prebiotic properties [77]. The well known skeletal benefits of VD are in part mediated by butyrate. Butyrate stimulates osteoblastic activity and down-regulates osteoporosis, possibly by stimulating the release of PTH [86,87]. The opposite is also true. Gut dysbiosis is linked with suboptimal VD status [88]. The gut microbiome regulates not only bone homeostasis and bone health but also many extraskeletal functions of VD, e.g., anti cancer, anti diabetes, anti hypertension, anti obesity, anti dementia, anti autoimmunity [89]. Dysbiosis compromises the absorption of D3, Ca, and Mg. The gut Firmicutes/Bacteroidetes ratio is negatively linked to a healthy gut microbiome with suppression of Firmicutes phylum bacteria upon supplementation with 25(OH) D or D3 [90,91]. VD deficiency also negatively impacts the gut microbiome, compromising B vitamin production.

While Mg is required for activation of B2,3,6,9,12, the gut microbiome may fill in for any shortfall [92]. Some intestinal bacteria can produce all eight B vitamins and up to 65% of human gut microorganisms can synthesize at least one type of B vitamin [93,94]. The few discrepant reports on the efficacy of D3 with respect to the gut microbiome may be due to:

1. Lack of baseline data indicating insufficiency/deficiency

2. Failure to property separate placebo and target groups by baseline

3. Less than 2-3 months between start of D3 supplementation and measurement of results

4. Insufficient D3 dosage

5. Failure to normalize for Ca:Mg as a confounding factor

6. Target group too small

2.5 Magnesium and the Gut Microbiome

Primary bile acids are metabolized by Mg dependent CYP450 (Cytochrome P450) enzymes and impact the gut microbiome [95,96]. Primary bile acids are metabolized to secondary bile acids by intestinal bacteria. Secondary bile acids, associated

with longevity, are also Mg dependent and require a healthy gut microbiome [97]. Both SCFAs and secondary bile acids independently exert a myriad of beneficial effects on host health [98]. Recently the vital role of aryl hydrocarbon receptors (AhRs) in aging, dementia, autoimmune disease, cancer, and ASCVD [99-101] has been recognized. AhR activity is dependent on its ligands. Kynurenines are AhR ligands that accelerate aging and neurodegeneration, while butyrates and indoles of intestinal bacterial origin are AhR ligands that oppose this and promote longevity. Indoles are longevity agents produced by gut microbiota. Their healthful benefits depend on the aryl hydrocarbon receptor (AhR) [99,102]. Not only are indoles and butyrate AhR ligands, but their subsequent AhR activation induces Mg dependent cytochrome CYP450 enzymes that facilitate gut absorption of indoles [103,104]. All CYP450 enzymes are Mg dependent [91]. Like Ozempic, butyrate is also a GLP-1 (glucagon-like peptide) agonist, albeit a natural one. Both are longevity indicators. Probiotics, rich in butyrogenic bacteria, are also associated with longevity and Mg enhances their efficacy [105-107]. Butyrate produced by gut bacteria may via vagal afferents increase HRV, linked to longevity. Gut microbiota cannot produce SCFAs in the absence of Mg, but they can produce tryptophan, another longevity agent. Candida yeast and hyphal forms produce their own form of IDO that competes with host IDO [108-112]. It enhances degradation of tryptophan to kynurenine (K/T) and accelerates the kynurenine pathway, increasing K/T, especially under the direction of interferon-gamma [113]. Any shortfall in Mg can increase subsequent neurotoxic metabolites and decrease NAD+ production [2]. The K/T ratio is negatively linked to the health of the gut microbiome [113]. Lactobacilli and Bifidobacteria boost plasma tryptophan levels [111]. They also produce lactate, which can crossfeed butyrogenic bacteria [114]. Gut microbes provide most of our circulating tryptophan. However, in order to absorb it intestinal epithelial cells rely on B0AT, a neutral or nonpolar amino acid transporter, that works in concert with ACE2 receptors [115]. This puts those with Covid-19, long Covid, or Candida overgrowth at risk for gut dysbiosis, compromising the enormous benefits of VD and Mg.

2.6 Therapeutic Interventions (Figure 7)

2.7 Probiotics, e.g., yogurt, alone are insufficient, if diet is suboptimal. The "good" bacteria must be fed and require fiber or indigestible carbohydrates, i.e., prebiotic, e.g.,

d-mannose. Butyrate is a commercially available postbiotic (produced by the "good" bacteria) that mimics the actions of Ozempic.

2.8 Target 2.0 for iCa:iMg or a range of 1.7 to 2.8 Ca:Mg in mmoles/L. If elevated, lower dietary Ca, e.g., eliminate dairy products, first and then increase dietary Mg, e.g., nuts, seeds, leafy greens, avocados. The silent damage overtime due to an elevated or depressed ratio escapes detection and the laxative effect of Mg facilitates this. Ca:Mg in the "healthy" has well exceeded 3.0 for at least two decades. Chronic latent Mg deficiency remains hidden, when the lower limit of normal persists at 0.75 mmol/L. The 1.7-2.8 ratio range represents a proposed Ca:Mg RDA, depending on

diet, BMI and age [37].

2.9 After improving Ca:Mg supplement D3 to attain a serum level of at least 50 ng/mL (125 mM) 25(OH)D (Figure 7) [116,117].

2.10 Take supplemental Mg with pyridoxal phosphate, the active form of B6, and perhaps D3. Mg is required for the hydroxylation of D3 in the liver (storage form). Taking pyridoxal phosphate concomitantly with Mg can enhance absorption and availability of Mg [118,119]. Not only does pyridoxal phosphate enhance cellular uptake of Mg but Mg enhances that of pyridoxal phosphate [120]. Several studies have challenged this [121,122]. But both studies employed the inactive form-pyridoxine. Lysine also enhances the absorbability of Mg.

2.11 Avoid simultaneous Ca and Mg intake. Although CaSRs are primarily found in the parathyroid gland and the kidney, they are also present in many other organs, including the alimentary canal [123].

2.12 Avoid simultaneous processed food/soft drinks and Mg intake. The former contain phosphates, which bind both Ca and

Mg in that order, limiting absorption. This may in part explain the global shortfall in Ca and Mg absorption.

2.13 Exercise induced elevation of lactate may enhance serum butyrate. Lactate may permeate intestinal endothelial and epithelial cells into the alimentary canal, where it can crossfeed butyrogenic bacteria [114].

2.14 Pay close attention to proper hydration. Dehydration triggers release of aldosterone, which increases renal reabsorption of Na+ and urinary excretion of Mg++ and K+. Cortisol possesses properties similar to aldosterone and can to a lesser extent trigger this same cationic exchange. Stress induced cortisol can lead to Mg deficiency, while Mg deficiency in turn enhances the body's susceptibility to stress [124].

2.15 Increase VD intake with age and increasing morbidity.

2.16 Replenish water soluble B vitamins that require Mg for activation and are required for synthesis of 7-dehydrocholesterol from acetate, enabling solar conversion to D3, (Figure 1).



Figure 7: This illustrates a proposed flowchart for deficiencies of VD, Ca, and Mg in various racial, ethnic, cultural, and economic sub groups, many of which overlap. Flow from a suboptimal gut microbiome at the bottom in red through a suggested therapeutic approach in yellow to an optimal gut microbiome and balanced Ca:Mg (green) is indicated. Rx=therapy, Sup=supplementation

3. Conclusion

Mg may be to VD what the gut microbiome is to general health. One is indispensable to the other. This review presents the benefits of both to gut microbiome dependent longevity agents. Many indirect benefits of this partnership are not included, e.g., maintenance of intestinal integrity [107,125]. VD and Mg are inextricably entwined and work hand in glove. Thirty ng/ mL of 25(OH)D and 0.75 mg/dL of Mg are insufficient. Mg is indispensable for not only the synthesis of VD's solar substrate (D3) and the storage/active forms of VD from D3 but also the synthesis and secretion of PTH. A healthy gut microbiome is also required to fully realize the benefits of both. Furthermore, there are many other gut microbiome related micronutrients and vitamins that depend on VD and Mg for their healthful effects and vice versa. Indeed there are myriad extraskeletal and extraintestinal benefits of both. The interdependencies are both legion and complex. Many have only recently been discovered and much remains hidden. The overlooked but critical role of the Ca:Mg ratio is at odds with their RDAs and compromises the efficacy of supplemental D3. The ratio can be easily calculated from serum Ca and Mg in the non hospitalized.

This review is largely hypothetical and highlights the relevant physiology, while overlooking the massive role of genes. The contributions of the gut microbiome are massively underappreciated. The increasingly sedentary Western lifestyle, the deteriorating quality of food, the abundant use of antibiotics and certain other medications, e.g., proton pump inhibitors and some diuretics, have conspired to challenge the quality of our gut microbiome. Not surprisingly the benefits of fecal microbiota transplantation (FMT) have rapidly expanded from its initial treatment for fulminant pseudomembranous colitis due to Clostridium difficile to an emerging tool for alleviating diseases related to a problematic gut microbiome. Gut microbiome manipulation and FMT may eventually become the frontline therapeutic approach to all diseases from cancer and dementia to Lyme disease [126]. VD, Ca, and Mg are majority players in this manipulation along with prebiotics + probiotics (synbiotics) and postbiotics. VD efficacy is tightly linked to Ca:Mg and VD studies that adjust for race, ethnicity, culture, socioeconomic status, and diet may even reveal greater efficacy. Unfortunately many of these conditions overlap in complex ways and difficulties in evaluating Mg status have made this a challenging proposition. This speculative view of VD efficacy is supported by many recent Covid-19 driven reports and relevant laboratory data. However, robust analysis of clinical correlations awaits.

References

- Fantini, C., Corinaldesi, C., Lenzi, A., Migliaccio, S., & Crescioli, C. (2023). Vitamin D as a Shield against Aging. *International Journal of Molecular Sciences*, 24(5), 4546
- 2. Chambers, P. (2024). Magnesium and Longevity. Qeios.
- Badal, V. D., Vaccariello, E. D., Murray, E. R., Yu, K. E., Knight, R., Jeste, D. V., & Nguyen, T. T. (2020). The Gut Microbiome, Aging, and Longevity: A Systematic Review. *Nutrients*, 12(12), 3759.

- Bidell, M.R., Hobbs, A.L.V., Lodise, T.P. (2022). Gut microbiome health and dysbiosis: A clinical primer. *Pharmacotherapy*, 42(11), 849-857.
- Martinez, J.E., Kahana, D.D., Ghuman, S., Wilson, H.P., Wilson, J., Kim, S.C.J., et al. (2021). Unhealthy Lifestyle and Gut Dysbiosis: A Better Understanding of the Effects of Poor Diet and Nicotine on the Intestinal Microbiome. *Front Endocrinol, 12,* 667066.
- 6. Norman, D.A., Fordtran, J.S., Brinkley, L.J., Zerwekh, J.E., Nicar, M.J., Strowig, S.M, et al. (1981). Jejunal and ileal adaptation to alterations in dietary calcium: changes in calcium and magnesium absorption and pathogenetic role of parathyroid hormone and 1,25-dihydroxyvitamin D. *J Clin Invest*, *67*(6), 1599-603.
- Mahaffee, D.D., Cooper, C.W., Ramp, W.K., Ontjes, D.A. (1982). Magnesium promotes both parathyroid hormone secretion and adenosine 3',5'-monophosphate production in rat parathyroid tissues and reverses the inhibitory effects of calcium on adenylate cyclase. *Endocrinology*, 110(2), 487-95.
- 8. Brown, E.M., Hurwitz, S., Aurbach, G.D. (1977). Betaadrenergic stimulation of cyclic AMP content and parathyroid hormone release from isolated bovine parathyroid cells. *Endocrinology*, *100*(6), 1696-702.
- Mostafa, W.Z., Hegazy, R.A. (2015). Vitamin D and the skin: Focus on a complex relationship: A review. J Adv Res, 6(6), 793-804.
- Sankalya, S. Ambagaspitiya., Gayan, A. Appuhamillage., Sunil, J. Wimalawansa. (2025). Impact of Vitamin D on Skin Aging, and Age-Related Dermatological Conditions. *Front. Biosci. (Landmark Ed)*, 30(1), 25463
- 11. https://vitamindstopscovid.info/02-intracrine/
- Quraishi, S.A., Bittner, E.A., Blum, L., Hutter, M.M., Camargo, C.A. (2014). Association Between Preoperative 25-Hydroxyvitamin D Level and Hospital-Acquired Infections Following Roux-en-Y Gastric Bypass Surgery. *JAMA Surg*, 149(2), 112–118.
- 13. Israel, A., Cicurel, A., Feldhamer, I., et al. (2020). The link between vitamin D deficiency and Covid-19 in a large population. *medRxiv*, 2020.
- Veugelers, P.J., Ekwaru, J.P. (2014). A statistical error in the estimation of the recommended dietary allowance for vitamin D. Nutrients, 6, 4472–4475.
- 15. Chambers, P. (2022). Comment on Huțanu et al. Low serum vitamin D in COVID-19 patients is not related to inflammatory markers and patients' outcomes—A single-center experience and a brief review of the literature. *Nutrients, 14*(1998).
- 16. Sempos, C. T., Looker, A. C., Durazo-Arvizu, R. A., Yetley, E. A., Chaudhary-Webb, M., Maw, K. L., et al. (2016). The vitamin D status of the US population from 1988 to 2010 using standardized serum concentrations of 25-hydroxyvitamin D shows recent modest increases. *American Journal of Clinical Nutrition, 104*(2), 454-461.
- Subramanian, A., Burrowes, H. B., Rumph, J. T., Wilkerson, J., Jackson, C. L., & Jukic, A. M. Z. (2024). Vitamin D levels in the United States: Temporal trends (2011-2018) and contemporary associations with sociodemographic

characteristics (2017-2018). Nutrients, 16(19), 3414.

- Durlach, J. (1989). Recommended dietary amounts of magnesium: Mg RDA. *Magnesium Research*, 2(3), 195-203.
- Costello, R. B., Rosanoff, A., Dai, Q., Saldanha, L. G., & Potischman, N. A. (2021). Perspective: Characterization of dietary supplements containing calcium and magnesium and their respective ratio—Is a rising ratio a cause for concern? *Advances in Nutrition*, 12(2), 291-297.
- 20. Razzaque, M. S. (2018). Magnesium: Are we consuming enough? *Nutrients*, 10(12), 1863.
- 21. Elin, R.J. (2010). Assessment of magnesium status for diagnosis and therapy. *Magnes Res*, 23(4), S194-8.
- 22. Micke, O., Vormann, J., Kraus, A., Kisters, K. (2021). Serum Magnesium: Time for a Standardized and Evidence-Based Reference Range. *Magnetic Resonance*, *34*, 84-89.
- Rosanoff, A., West, C., Elin, R.J., Micke, O., Baniasadi, S., Barbagallo, M., et al. (2022). MaGNet Global Magnesium Project (MaGNet). Recommendation on an updated standardization of serum magnesium reference ranges. *Eur J Nutr, 61*(7), 3697-3706.
- Weiss, D., Brunk, D.K., Goodman, D.A. (2018). Scottsdale Magnesium Study: Absorption, Cellular Uptake, and Clinical Effectiveness of a Timed-Release Magnesium Supplement in a Standard Adult Clinical Population. *J Am Coll Nutr*, 37(4), 316-327.
- 25. Rosanoff, A., Weaver, C.M., Rude, R.K. (2012). Suboptimal magnesium status in the United States: are the health consequences underestimated? *Nutr Rev, 70*, 153-64.
- 26. Gagliano, V., Schäffeler, F., Del Giorno, R., Bianchetti, M., Carvajal Canarte, CF., Caballero Regueira, J.J., et al. (2022). Does Ionized Magnesium Offer a Different Perspective Exploring the Association between Magnesemia and Targeted Cardiovascular Risk Factors? *Journal of Clinical Medicine*, 11(14), 4015.
- Sava, L., Pillai, S., More, U., Sontakke, A. (2005). Serum calcium measurement: Total versus free (ionized) calcium. *Indian J Clin Biochem*, 20(2), 158-61.
- Quinn, S.J., Thomsen, A.R., Egbuna, O., Pang, J., Baxi K., Goltzman, D., et al. (2013). CaSR-mediated interactions between calcium and magnesium homeostasis in mice. *Am J Physiol Endocrinol Metab*, 304(7), E724-33.
- 29. Dai, Q., Zhu, X., Manson, J.E., Song, Y., Li, X., Franke, A., et al. (2018) Magnesium Status and Supplementation Influence Vitamin D Status and Metabolism: Results from a Randomized Trial. *The American Journal of Clinical Nutrition, 108*, 1249-1258.
- Cheung, M.M., DeLuccia, R., Ramadoss, R.K., Aljahdali, A., Volpe, S.L., Shewokis, P.A., et al. (2019). Low dietary magnesium intake alters vitamin D-parathyroid hormone relationship in adults who are overweight or obese. *Nutr Res*, 69, 82-93.
- Dai, Q., Shu, X.O., Deng, X., Xiang, Y.B., Li, H., Yang, G., et al. (2013). Modifying effect of calcium/magnesium intake ratio and mortality: a population-based cohort study. *BMJ Open*, 3(2), e002111.

- 32. Altura, B.M. (1994). Introduction: importance of Mg in physiology and medicine and the need for ion selective electrodes. *Scand J Clin Lab Invest Suppl, 217*, 5-9.
- 33. https://www.labcorp.com/tests/080283/magnesium-rbc
- Areco, V. A., Kohan, R., Talamoni, G., Tolosa de Talamoni, N. G., & Peralta López, M. E. (2020). Intestinal Ca2+ absorption revisited: A molecular and clinical approach. World Journal of Gastroenterology, 26(24), 3344-3364.
- 35. Labcorp reference range for ionized serum Ca
- Han, C., Shin, A., Lee, J. et al. (2015). Dietary calcium intake and the risk of colorectal cancer: a case control study. *BMC Cancer 15*, 966.
- Dai, Q., Motley, S. S., Smith, J. A. Jr., Concepcion, R., Barocas, D., Byerly, S., & Fowke, J. H. (2011). Blood magnesium, and the interaction with calcium, on the risk of high-grade prostate cancer. *PLOS One*, *6*(4), e18237.
- Zhao, J., Giri, A., Zhu, X. et al. (2019). Calcium: magnesium intake ratio and colorectal carcinogenesis, results from the prostate, lung, colorectal, and ovarian cancer screening trial. *Br J Cancer 121*, 796–804 (2019).
- Zhang, H., Cao, Y., Song, P., Man, Q., Mao, D., Hu, Y., & Yang, L. (2021). Suggested reference ranges of blood Mg and Ca level in childbearing women of China: Analysis of China Adult Chronic Disease and Nutrition Surveillance (2015). *Nutrients*, 13(9), 3287.
- Yang, J., Cao, Y., Shan, X., Zhang, H., Feng, J., Lu, J., & et al. (2023). The magnesium status and suggested reference ranges of plasma magnesium, calcium, and calcium/magnesium ratio in Chinese adults over 45 years old. *Nutrients*, 15(4), 886.
- 41. Rosanoff, A., Dai, Q., & Shapses, S. A. (2016). Essential nutrient interactions: Does low or suboptimal magnesium status interact with vitamin D and/or calcium status? *Advances in Nutrition*, *7*(1), 25-43
- Ginde, A. A., Wolfe, P., Camargo, C. A. Jr., & Schwartz, R. S. (2012). Defining vitamin D status by secondary hyperparathyroidism in the U.S. population. *Journal of Endocrinological Investigation*, 35(1), 42-48
- 43. Hyung-Suk, Yoon., Xiao Ou, Shu., Hui, Cai., Wei, Zheng., William, J. Blot., Qiuyin, Cai. (2021). Abstract 851: Associations of dietary calcium and magnesium intakes with lung cancer risk among low-income Americans: Results from the Southern Community Cohort Study. *Cancer Res*, 81, 851.
- 44. Huang, J.H., Tsai, L.C., Chang, Y.C. et al. (2014). High or low calcium intake increases cardiovascular disease risks in older patients with type 2 diabetes. *Cardiovasc Diabetol 13*, 120.
- Takata, Y., Yang, J. J., Yu, D., Smith-Warner, S. A., Blot, W. J., White, E., & et al. (2023). Calcium intake and lung cancer risk: A pooled analysis of 12 prospective cohort studies. *The Journal of Nutrition*, 153(7), 2051-2060.
- Hongwen, T., Wei, G., Quanwei, H., Mengjun, B., & Yingjiu, J. (2024). Research progress on the relationship between vitamin D and lung cancer. *Journal of Clinical Medicine Surgery*, 4(1), 1148.
- 47. Haider, F., Ghafoor, H., Hassan, O. F., Farooqui, K., Bel Khair, A. O. M., & Shoaib, F. (2023). Vitamin D and cardiovascular

diseases: An update. Cureus, 15(11), e49734.

- 48. Shah, S. C., Dai, Q., Zhu, X., Peek, R. M. Jr., Roumie, C., & Shrubsole, M. J. (2020). Associations between calcium and magnesium intake and the risk of incident oesophageal cancer: An analysis of the NIH-AARP Diet and Health Study prospective cohort. *British Journal of Cancer*, 122(12), 1857-1864.
- 49. Batai, K., Cui, Z., Arora, A., Shah-Williams, E., Hernandez, W., Ruden, M., & et al. (2021). Genetic loci associated with skin pigmentation in African Americans and their effects on vitamin D deficiency. *PLOS Genetics*, 17(2), e1009319.
- Scully, H., Laird, E., Healy, M., Crowley, V., Walsh, J. B., & McCarroll, K. (2022). Socioeconomic status predicts vitamin D status in a large cohort of Irish children. *Proceedings of the Nutrition Society, 81*(OCE4), E87.
- 51. Tønnesen, R., Hovind, P. H., & Jensen, L. T. (2016). Determinants of vitamin D status in young adults: Influence of lifestyle, sociodemographic and anthropometric factors. *BMC Public Health*, 16, 385
- 52. Hussein, D. A., et al. (2022). Pattern of vitamin D deficiency in a Middle Eastern population: A cross-sectional study. International Journal of Functional Nutrition.
- 53. Darling, A.L., et al. (2020) Very High Prevalence of 25-hydroxyvitamin D Deficiency in 6433 UK South Asian adults: analysis of the UK Biobank Cohort. British Journal of Nutrition.
- 54. Chan, Chin Yi., et al. (2018). Attitude of Asians to Calcium and Vitamin D Rich Foods and Supplements: A Systematic Review. Sains Malaysiana (2018).
- 55. NHANES data for nutrient intake by gender and age from 2001 through Aug 2023
- 56. Deng, X., Song, Y., Manson, J.E. et al. (2013). Magnesium, vitamin D status and mortality: results from US National Health and Nutrition Examination Survey (NHANES) 2001 to 2006 and NHANES III. *BMC Med*, 11, 187 (2013).
- 57. Nouri-Majd, S., Ebrahimzadeh, A., Mousavi, S. M., Zargarzadeh, N., Eslami, M., Santos, H. O., et al. (2022). Higher intake of dietary magnesium is inversely associated with COVID-19 severity and symptoms in hospitalized patients: A cross-sectional study. *Frontiers in Nutrition*, 9, 873162.
- Ginde, A. A., Brower, R. G., Caterino, J. M., Finck, L., Banner-Goodspeed, V. M., Grissom, C. K., et al. (2019). Early high-dose vitamin D3 for critically ill, vitamin D-deficient patients. *The New England Journal of Medicine*, 381(26), 2529-2540.
- Rooney, M. R., Fang, M., Ogurtsova, K., Ozkan, B., Echouffo-Tcheugui, J. B., Boyko, E. J., Magliano, D. J., & Selvin, E. (2023). Global prevalence of prediabetes. *Diabetes Care*, 46(7), 1388-1394.
- 60. Fahed, M., Abou Jaoudeh, M. G., Merhi, S., et al. (2020). Evaluation of risk factors for insulin resistance: A crosssectional study among employees at a private university in Lebanon. *BMC Endocrine Disorders, 20*, 85.
- 61. Rahim, N. E., Flood, D., Marcus, M. E., Theilmann, M., Aung, T. N., Agoudavi, K., et al. (2023). Diabetes risk and

provision of diabetes prevention activities in 44 low-income and middle-income countries: A cross-sectional analysis of nationally representative, individual-level survey data. *The Lancet Global Health*, *11*(10), e1576-e1586.

- 62. Dubey, P., Thakur, V., & Chattopadhyay, M. (2020). Role of minerals and trace elements in diabetes and insulin resistance. *Nutrients, 12*(6), 1864.
- Liu, H., Li, N., Jin, M., Miao, X., Zhang, X., & Zhong, W. (2020). Magnesium supplementation enhances insulin sensitivity and decreases insulin resistance in diabetic rats. *Iranian Journal of Basic Medical Sciences*, 23(8), 990-998.
- 64. Kostov, K. (2019). Effects of magnesium deficiency on mechanisms of insulin resistance in type 2 diabetes: Focusing on the processes of insulin secretion and signaling. *International Journal of Molecular Sciences*, 20(6), 1351.
- 65. Kherad, Z., Yazdanpanah, S., Saadat, F., Pakshir, K., & Zomorodian, K. (2023). Vitamin D3: A promising antifungal and antibiofilm agent against Candida species. *Current Medical Mycology*, 9(2), 17-22.
- Lei, J., Xiao, W., Zhang, J., Liu, F., Xin, C., Zhou, B., et al. (2022). Antifungal activity of vitamin D3 against Candida albicans in vitro and in vivo. *Microbiological Research*, 265, 127200.
- 67. Jawhara, S. (2022). How gut bacterial dysbiosis can promote Candida albicans overgrowth during colonic inflammation. *Microorganisms*, 10(5), 1014.
- 68. Thomas, R.L., Jiang, L., Adams, J.S. et al. (2020). Vitamin D metabolites and the gut microbiome in older men. *Nat Commun*, *11*, 5997.
- Tangestani, H., Boroujeni, H. K., Djafarian, K., Emamat, H., & Shab-Bidar, S. (2021). Vitamin D and the gut microbiota: A narrative literature review. *Clinical Nutrition Research*, 10(3), 181-191.
- 70. Singh, P., Rawat, A., Alwakeel, M. et al. (2020). The potential role of vitamin D supplementation as a gut microbiota modifier in healthy individuals. *Sci Rep, 10*, 21641.
- 71. Evangelos, G., et al. (2024). Vitamin D regulates microbiomedependent cancer immunity. *Science*, *384*, 428-437.
- 72. Yamamoto, E. A., & Jørgensen, T. N. (2020). Relationships between vitamin D, gut microbiome, and systemic autoimmunity. *Frontiers in Immunology*, *10*, 3141.
- Tabatabaeizadeh, S. E., Tafazoli, N., Ferns, G. A., Avan, A., & Ghayour-Mobarhan, M. (2018). Vitamin D, the gut microbiome, and inflammatory bowel disease. *Journal of Research in Medical Sciences*, 23(1), 75.
- 74. Kanhere, M., He, J., Chassaing, B., Ziegler, T. R., Alvarez, J. A., Ivie, E. A., et al. (2018). Bolus weekly vitamin D3 supplementation impacts gut and airway microbiota in adults with cystic fibrosis: A double-blind, randomized, placebo-controlled clinical trial. *The Journal of Clinical Endocrinology & Metabolism, 103*(2), 564–574.
- Cantarel, B. L., Waubant, E., Chehoud, C., Kuczynski, J., DeSantis, T. Z., Warrington, J., Venkatesan, A., Fraser, C. M., & Mowry, E. M. (2015). Gut microbiota in multiple sclerosis: Possible influence of immunomodulators. *Journal*

of Investigative Medicine, 63(5), 729-734.

- 76. Velizarova, M., Yanachkova, V., Boneva, T., Giragosyan, S., Mihaleva, I., Andreeva-Gateva, P., et al. (2023). Relationship between vitamin D status and microbiome changes in Bulgarian patients with type 2 diabetes mellitus. *Biotechnology* & *Biotechnological Equipment*, 37(1).
- 77. Daley, D. K., & Myrie, S. B. (2024). Diabetes and vitamin D: The effect of insulin sensitivity and gut microbial health. *Advances in Food and Nutrition Research*, 109, 160-184.
- Breuling, M., Tomeva, E., Ivanovic, N., & Haslberger, A. (2024). Butyrate- and beta-hydroxybutyrate-mediated effects of interventions with pro- and prebiotics, fasting, and caloric restrictions on depression: A systematic review and metaanalysis. *Life*, 14(7), 787.
- 79. Wang, C., Zheng, D., & Weng, F. et al. (2022). Sodium butyrate ameliorates the cognitive impairment of Alzheimer's disease by regulating the metabolism of astrocytes. *Psychopharmacology, 239,* 215–227.
- Chen, J., Zhao, K.-N., & Vitetta, L. (2019). Effects of intestinal microbial–elaborated butyrate on oncogenic signaling pathways. *Nutrients*, 11(5), 1026.
- Zhang, Y., Tao, Y., Gu, Y., & Ma, Q. (2023). Butyrate facilitates immune clearance of colorectal cancer cells by suppressing STAT1-mediated PD-L1 expression. Clinics (São Paulo, Brazil), 78, 100303.
- Coppola, S., Avagliano, C., Calignano, A., & Berni Canani, R. (2021). The protective role of butyrate against obesity and obesity-related diseases. *Molecules*, 26(3), 682.
- Blaak, E. E., Canfora, E. E., Theis, S., Frost, G., Groen, A. K., Mithieux, G., et al. (2020). Short-chain fatty acids in human gut and metabolic health. *Beneficial Microbes*, 11(5), 411-455.
- Facchin, S., Bertin, L., Bonazzi, E., Lorenzon, G., De Barba, C., Barberio, B., Zingone, F., Maniero, D., Scarpa, M., Ruffolo, C., et al. (2024). Short-chain fatty acids and human health: From metabolic pathways to current therapeutic implications. *Life*, 14(5), 559.
- 85. Charoenngam, N., Shirvani, A., Kalajian, T. A., Song, A., & Holick, M. F. (2020). The effect of various doses of oral vitamin D3 supplementation on gut microbiota in healthy adults: A randomized, double-blinded, dose-response study. *Anticancer Research*, 40(1), 551-556.
- Cooney, O. D., Nagareddy, P. R., Murphy, A. J., & Lee, M. K. S. (2021). Healthy gut, healthy bones: Targeting the gut microbiome to promote bone health. *Frontiers in Endocrinology*, 11, 620466.
- Pacifici, R. (2021). Role of gut microbiota in the skeletal response to PTH. *The Journal of Clinical Endocrinology & Metabolism*, 106(3), 636–645.
- Su, D., Nie, Y., Zhu, A., Chen, Z., Wu, P., Zhang, L., et al. (2016). Vitamin D signaling through induction of Paneth cell defensins maintains gut microbiota and improves metabolic disorders and hepatic steatosis in animal models. *Frontiers in Physiology*, 7, 498.
- 89. Hansdah, K., & Lui, J. C. (2024). Emerging insights into the endocrine regulation of bone homeostasis by gut microbiome.

Journal of the Endocrine Society, 8(8), bvae117.

- Rinninella, E., Raoul, P., Cintoni, M., Franceschi, F., Miggiano, G. A. D., Gasbarrini, A., & Mele, M. C. (2019). What is the healthy gut microbiota composition? A changing ecosystem across age, environment, diet, and diseases. *Microorganisms*, 7(1), 14.
- Bellerba, F., Muzio, V., Gnagnarella, P., Facciotti, F., Chiocca, S., Bossi, P., Cortinovis, D., Chiaradonna, F., Serrano, D., Raimondi, S., et al. (2021). The association between vitamin D and gut microbiota: A systematic review of human studies. *Nutrients*, 13(10), 3378.
- 92. Gominak, S. C. (2016). Vitamin D deficiency changes the intestinal microbiome, reducing B vitamin production in the gut. The resulting lack of pantothenic acid adversely affects the immune system, producing a "pro-inflammatory" state associated with atherosclerosis and autoimmunity. *Medical Hypotheses, 94*, 103-107.
- 93. Wibowo, S., & Pramadhani, A. (2024). Vitamin B, role of gut microbiota and gut health. IntechOpen.
- Nysten, J., & Van Dijck, P. (2023). Can we microbe-manage our vitamin acquisition for better health? *PLoS Pathogens*, 19(5).
- 95. Mansmann, H. C. (1994). Consider magnesium homeostasis: III: Cytochrome P450 enzymes and drug toxicity. *Applied Immunohistochemistry & Molecular Morphology*, *8*, 7-28.
- 96. Collins, S. L., Stine, J. G., Bisanz, J. E., et al. (2023). Bile acids and the gut microbiota: Metabolic interactions and impacts on disease. *Nature Reviews Microbiology*, *21*, 236–247.
- 97. Ji, S., Pan, Y., Zhu, L., Tan, J., Tang, S., Yang, Q., Zhang, Z., Lou, D., & Wang, B. (2021). A novel 7α-hydroxysteroid dehydrogenase: Magnesium ion significantly enhances its activity and thermostability. *International Journal of Biological Macromolecules*, 177, 111–118.
- 98. Kim, D. M., Liu, J., Whitmore, M. A., et al. (2024). Two intestinal microbiota-derived metabolites, deoxycholic acid and butyrate, synergize to enhance host defense peptide synthesis and alleviate necrotic enteritis. *Journal of Animal Science and Biotechnology*, 15, 29.
- 99. Ojo, E. S., & Tischkau, S. A. (2021). The role of AhR in the hallmarks of brain aging: Friend and foe. *Cells*, *10*(10), 2729.
- 100. Wang, Z., Snyder, M., Kenison, J. E., Yang, K., Lara, B., Lydell, E., et al. (2020). How the AhR became important in cancer: The role of chronically active AhR in cancer aggression. *International Journal of Molecular Sciences*, 22(1), 387.
- 101.Zhu, K., Meng, Q., Zhang, Z., Yi, T., He, Y., Zheng, J., et al. (2019). Aryl hydrocarbon receptor pathway: Role, regulation, and intervention in atherosclerosis therapy (Review). *Molecular Medicine Reports*, 20(6), 4763–4773.
- 102.Koper, J. E. B., Kortekaas, M., Loonen, L. M. P., Huang, Z., Wells, J. M., Gill, C. I. R., et al. (2020). Aryl hydrocarbon receptor activation during in vitro and in vivo digestion of raw and cooked broccoli (Brassica oleracea var. italica). *Food & Function*, 11(5), 4026–4037.
- 103. Marinelli, L., Martin-Gallausiaux, C., Bourhis, J. M., et al.

(2019). Identification of the novel role of butyrate as AhR ligand in human intestinal epithelial cells. *Scientific Reports*, *9*, 643.

- 104. Li, X., Zhang, B., Hu, Y., & Zhao, Y. (2021). New insights into gut-bacteria-derived indole and its derivatives in intestinal and liver diseases. *Frontiers in Pharmacology*, 12, 769501.
- 105.Gribble, F. M., & Reimann, F. (2021). Metabolic messengers: Glucagon-like peptide 1. Nature Metabolism, 3, 142–148.
- 106. Chaudhary, P., Kathuria, D., Suri, S., Bahndral, A., & Kanthi Naveen, A. (2023). Probiotics—Its functions and influence on the aging process: A comprehensive review. *Food Bioscience*, 52, 102389.
- 107. Mahboobi, S., Ghasvarian, M., Ghaem, H., Alipour, H., Alipour, S., & Eftekhari, M. H. (2022). Effects of probiotic and magnesium co-supplementation on mood, cognition, intestinal barrier function, and inflammation in individuals with obesity and depressed mood: A randomized, doubleblind, placebo-controlled clinical trial. *Frontiers in Nutrition*, 9, 1018357.
- 108. Seefeldt, J. M., Homilius, C., Hansen, J., Lassen, T. R., Jespersen, N. R., Jensen, R. V., et al. (2024). Short-chain fatty acid butyrate is an inotropic agent with vasorelaxant and cardioprotective properties. *Journal of the American Heart Association: Cardiovascular and Cerebrovascular Disease*, 13.
- 109. Yu, Z., Han, J., Chen, H., Wang, Y., Zhou, L., Wang, M., et al. (2021). Oral supplementation with butyrate improves myocardial ischemia/reperfusion injury via a gut-brain neural circuit. *Frontiers in Cardiovascular Medicine*, *8*, 718674.
- 110. Sasaki, H., Hayashi, K., Imamura, M., Hirota, Y., Hosoki, H., Nitta, L., et al. (2023). Combined resistant dextrin and low-dose magnesium oxide administration increases shortchain fatty acid and lactic acid production by gut microbiota. *Journal of Nutritional Biochemistry*, 120, 109420.
- 111. Hou, Y., Li, J., & Ying, S. (2023). Tryptophan metabolism and gut microbiota: A novel regulatory axis integrating the microbiome, immunity, and cancer. *Metabolites*, 13(11), 1166.
- 112. Kherad, Z., Yazdanpanah, S., Saadat, F., Pakshir, K., & Zomorodian, K. (2023). Vitamin D3: A promising antifungal and antibiofilm agent against Candida species. *Current Medical Mycology*, 9(2), 17–22.
- 113. Campbell, B. M., Charych, E., Lee, A. W., & Möller, T. (2014). Kynurenines in CNS disease: Regulation by inflammatory cytokines. *Frontiers in Neuroscience*, *8*, 12.
- 114. Louis, P., Duncan, S. H., Sheridan, P. O., Walker, A. W., & Flint, H. J. (2022). Microbial lactate utilization and the stability of the gut microbiome. Gut Microbiome, 3, e3.

- 115. Li, J., Yan, Y., & Fu, Y. (2024). ACE2 mediates tryptophan alleviation on diarrhea by repairing intestine barrier involved mTOR pathway. *Cellular and Molecular Biology Letters, 29*, 90.
- 116. Wimalawansa, S. J. (2024). Physiology of vitamin D— Focusing on disease prevention. *Nutrients, 16*(11), 1666.
- 117. AlHewishel, M. A., Bahgat, M., Al Huwaiyshil, A., et al. (2020, June 29). 25(OH)D serum level in non-diabetic and type II diabetic patients: A cross-sectional study. *Cureus*, *12*(6), e8910.
- 118. Abraham, G. E., Schwartz, U. D., & Lubran, M. M. (1981). Effect of vitamin B-6 on plasma and red blood cell magnesium levels in premenopausal women. *Annals of Clinical & Laboratory Science*, 11(4), 333–336.
- 119. Boylan, L. M., & Spallholz, J. E. (1990). In vitro evidence for a relationship between magnesium and vitamin B-6. Magnesium Research, 3, 79–85. https://pubmed.ncbi.nlm.nih. gov/2133627
- 120.Planells, E., Lerma, A., Sánchez-Morito, N., Aranda, P., & Llopis, J. (1997). Effect of magnesium deficiency on vitamin B2 and B6 status in the rat. *Journal of the American College of Nutrition, 16*(4), 352–356.
- 121.Noah, L., Pickering, G., Dubray, C., Mazur, A., & Hitier, S., et al. (2020). Effect of vitamin B6 supplementation, in combination with magnesium, on severe stress and magnesium status: Secondary analysis from an RCT. *Proceedings of the Nutrition Society, 79*(OCE2), E491.
- 122.Pouteau, E., Kabir-Ahmadi, M., Noah, L., Mazur, A., & Dye, L., et al. (2018). Superiority of magnesium and vitamin B6 over magnesium alone on severe stress in healthy adults with low magnesemia: A randomized, single-blind clinical trial. *PLoS One*, 13(12), e0208454.
- 123.Ohsu, T., Amino, Y., Nagasaki, H., Yamanaka, T., Takeshita, S., Hatanaka, T., et al. (2010). Involvement of the calciumsensing receptor in human taste perception. *The Journal of Biological Chemistry*, 285(2), 1016–1022.
- 124.Pickering, G., Mazur, A., Trousselard, M., Bienkowski, P., Yaltsewa, N., Amessou, M., Noah, L., & Pouteau, E. (2020). Magnesium status and stress: The vicious circle concept revisited. *Nutrients*, *12*(12), 3672.
- 125. Akimbekov, N.S, Digel, I., Sherelkhan, D.K., Lutfor, A.B., Razzaque, M.S. (2020). Vitamin D and the Host-Gut Microbiome: A Brief Overview. *Acta Histochem Cytochem*, 26;53(3):33-42.
- 126. Morrissette, M, Pitt, N, González, A, Strandwitz, P, Caboni, M, Rebman, AW. (2020). A Distinct Microbiome Signature in Posttreatment Lyme Disease Patients mBio, 11 (5) e02310-20.

Copyright: ©2025 Patrick Chambers. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.